

UBC9 (H-81): sc-10759

BACKGROUND

UBC9 is a component of the ubiquitin-mediated proteolytic pathway, which targets proteins for degradation by the 26S proteasome, mediates endocytosis and directs protein subcellular localization. Ub and Ub-like molecules are systematically transferred from E2 conjugating enzymes to the targeted substrate by way of an E3 ubiquitin ligase. UBC9 functions as an E2 ubiquitin conjugating enzyme that preferentially associates with the ubiquitin homolog designated SUMO-1 or sentrin, a component of the sentrinization complex. Characteristic of the E2 family members, UBC9 contains a conserved cysteine residue that is required for the thio ester formation between Ub-like proteins and the E2 member, and it shares a conserved UBC domain. Substrates for UBC9 include transcription factors E12 and E47 and mitotic regulators Ran BP-2 and Ran GAP1, which indicates that UBC9 may regulate a variety of cellular processes including cell cycle progression and differentiation.

CHROMOSOMAL LOCATION

Genetic locus: UBE2I (human) mapping to 16p13.3; Ube2i (mouse) mapping to 17 A3.3.

SOURCE

UBC9 (H-81) is a rabbit polyclonal antibody raised against amino acids 1-81 of UBC9 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

UBC9 (H-81) is recommended for detection of UBC9 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

UBC9 (H-81) is also recommended for detection of UBC9 in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for UBC9 siRNA (h): sc-36773, UBC9 siRNA (m): sc-36774, UBC9 shRNA Plasmid (h): sc-36773-SH, UBC9 shRNA Plasmid (m): sc-36774-SH, UBC9 shRNA (h) Lentiviral Particles: sc-36773-V and UBC9 shRNA (m) Lentiviral Particles: sc-36774-V.

Molecular Weight of UBC9: 18 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, Jurkat whole cell lysate: sc-2204 or U-937 cell lysate: sc-2239.


STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.


RESEARCH USE

For research use only, not for use in diagnostic procedures.

DATA



UBC9 (H-81): sc-10759. Western blot analysis of UBC9 expression in Jurkat whole cell lysate.



UBC9 (H-81): sc-10759. Western blot analysis of UBC9 expression in HeLa whole cell lysate.

SELECT PRODUCT CITATIONS

- Floyd, Z.E., et al. 2004. Control of peroxisome proliferator-activated receptor γ 2 stability and activity by SUMOylation. *Obes. Res.* 12: 921-928.
- Droescher, M., et al. 2011. Cytokine-induced paracrystals prolong the activity of signal transducers and activators of transcription (STAT) and provide a model for the regulation of protein solubility by small ubiquitin-like modifier (SUMO). *J. Biol. Chem.* 286: 18731-18746.
- Haghikia, A., et al. 2011. Signal transducer and activator of transcription 3-mediated regulation of miR-199a-5p links cardiomyocyte and endothelial cell function in the heart: a key role for ubiquitin-conjugating enzymes. *Eur. Heart J.* 32: 1287-1297.
- Qin, Y., et al. 2011. Ubc9 mediates nuclear localization and growth suppression of BRCA1 and BRCA1a proteins. *J. Cell. Physiol.* 226: 3355-3367.
- Suda, N., et al. 2011. Coactivation of SF-1-mediated transcription of steroidogenic enzymes by Ubc9 and PIAS1. *Endocrinology* 152: 2266-2277.
- Chu, Y., et al. 2011. SUMO E3 ligase activity of TRIM proteins. *Oncogene* 30: 1108-1116.
- Fuhs, S.R., et al. 2011. Caveolin-3 undergoes SUMOylation by the SUMO E3 ligase PIASy: sumoylation affects G-protein-coupled receptor desensitization. *J. Biol. Chem.* 286: 14830-14841.
- Liu, S.T., et al. 2012. A non-covalent interaction between small ubiquitin-like modifier-1 and Zac1 regulates Zac1 cellular functions. *Int. J. Biochem. Cell Biol.* 44: 547-555.



Try **UBC9 (C-12): sc-271057** or **UBC9 (50): sc-136245**, our highly recommended monoclonal alternatives to UBC9 (H-81). Also, for AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647 conjugates, see **UBC9 (C-12): sc-271057**.