

AKAP 79 (H-105): sc-10764

BACKGROUND

The type II cAMP-protein kinase (PKA) is a multifunctional kinase with a broad range of substrates. Specificity of PKA signaling is thought to be mediated by the compartmentalization of the kinase to specific sites within the cell. To maintain this specific localization, the R subunit (RII) of PKA interacts with specific RII-anchoring proteins. This family of proteins has been designated A-kinase anchoring proteins (AKAP). Members of this family, including MAP2 (microtubule-associated protein 2), neuronally expressed AKAP 79 and AKAP 150, and the DNA binding AKAP 95, display differential tissue specificity and localization. Evidence suggests that AKAP 79 and AKAP 150 are both capable of anchoring PKA to postsynaptic densities (PSD), which are a network of proteins located on the internal surfaces of excitatory synapses.

REFERENCES

1. Scott, J.D., et al. 1990. Type II regulatory subunit dimerization determines the subcellular localization of the cAMP-dependent protein kinase. *J. Biol. Chem.* 265: 21561-21566.
2. Carr, D.W., et al. 1992. Localization of the cAMP-dependent protein kinase to the postsynaptic densities by A-kinase anchoring proteins. Characterization of AKAP 79. *J. Biol. Chem.* 267: 16816-16823.
3. Coghlan, V.M., et al. 1993. A-kinase anchoring proteins: a key to selective activation of cAMP-responsive events? *Mol. Cell. Biochem.* 127: 309-319.
4. Coghlan, V.M., et al. 1994. Cloning and characterization of AKAP 95, a nuclear protein that associates with the regulatory subunit of type II cAMP-dependent protein kinase. *J. Biol. Chem.* 269: 7658-7665.
5. Coghlan, V.M., et al. 1995. Association of protein kinase A and protein phosphatase 2 β with a common anchoring protein. *Science* 267: 108-111.
6. Lester, L.B., et al. 1996. Cloning and characterization of a novel A-kinase anchoring protein. AKAP 220, association with testicular peroxisomes. *J. Biol. Chem.* 271: 9460-9465.

CHROMOSOMAL LOCATION

Genetic locus: AKAP5 (human) mapping to 14q21-q24.

SOURCE

AKAP 79 (H-105) is a rabbit polyclonal antibody raised against amino acids 181-285 of AKAP 79 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

AKAP 79 (H-105) is recommended for detection of AKAP 79 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for AKAP 79 siRNA (h): sc-29660, AKAP 79 shRNA Plasmid (h): sc-29660-SH and AKAP 79 shRNA (h) Lentiviral Particles: sc-29660-V.


Molecular Weight of AKAP 79: 79 kDa.

Positive Controls: IMR-32 cell lysate: sc-2409, H4 cell lysate: sc-2408 or JAR cell lysate: sc-2276.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



AKAP 79 (H-105): sc-10764. Western blot analysis of AKAP 79 expression in H4 (A) and JAR (B) whole cell lysates.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **AKAP 79 (D-9): sc-17772** or **AKAP 79 (22): sc-135826**, our highly recommended monoclonal alternatives to AKAP 79 (H-105).