

Med22 (T-15): sc-107739

BACKGROUND

Med22 is a subunit of the RNA polymerase II (Pol II) transcriptional mediator complex. The mediator complex is a coactivator involved in the regulated transcription of Pol II-dependent genes. Functioning as a bridge to convey information from gene-specific regulatory proteins to the basal Pol II transcription machinery, the mediator complex is recruited to promoter regions by directly interacting with regulatory proteins. The mediator complex also serves as a scaffold for the assembly of a functional pre-initiation complex with Pol II and other general transcription factors. Med22 (mediator complex subunit 22), also known as SURF5 (surfeit locus protein 5), is a ubiquitously expressed 200 amino acid nuclear protein that is one of several components of the mediator complex. There are two isoforms of Med22 that are produced as a result of alternative splicing events.

REFERENCES

- Garson, K., Duhig, T., Armes, N., Colombo, P. and Fried, M. 1995. Surf5: a gene in the tightly clustered mouse surfeit locus is highly conserved and transcribed divergently from the rpl7A (Surf3) gene. *Genomics* 30: 163-170.
- Garson, K., Duhig, T. and Fried, M. 1996. Tissue-specific processing of the Surf-5 and Surf-4 mRNAs. *Gene Expr.* 6: 209-218.
- Duhig, T., Ruhrberg, C., Mor, O. and Fried, M. 1998. The human surfeit locus. *Genomics* 52: 72-78.
- Angiolillo, A., Russo, G., Porcellini, A., Smaldone, S., D'Alessandro, F. and Pietropaolo, C. 2002. The human homologue of the mouse Surf5 gene encodes multiple alternatively spliced transcripts. *Gene* 284: 169-178.
- Martinez, E. 2002. Multi-protein complexes in eukaryotic gene transcription. *Plant Mol. Biol.* 50: 925-947.
- Esnault, C., Ghavi-Helm, Y., Brun, S., Soutourina, J., Van Berkum, N., Boschiero, C., Holstege, F. and Werner, M. 2008. Mediator-dependent recruitment of TFIID modules in preinitiation complex. *Mol. Cell* 31: 337-346.
- Online Mendelian Inheritance in Man, OMIM™. 2008. Johns Hopkins University, Baltimore, MD. MIM Number: 185641. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: MED22 (human) mapping to 9q34.2; Med22 (mouse) mapping to 2 A3.

SOURCE

Med22 (T-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Med22 of human origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-107739 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Med22 (T-15) is recommended for detection of Med22 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Med22 siRNA (h): sc-92912, Med22 siRNA (m): sc-149351, Med22 shRNA Plasmid (h): sc-92912-SH, Med22 shRNA Plasmid (m): sc-149351-SH, Med22 shRNA (h) Lentiviral Particles: sc-92912-V and Med22 shRNA (m) Lentiviral Particles: sc-149351-V.

Molecular Weight of Med22: 22 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfaction
Guaranteed

Try **Med22 (D-10): sc-393738**, our highly recommended monoclonal alternative to Med22 (T-15).