



# MIS12 (C-13): sc-107750

## BACKGROUND

Chromosome segregation requires assembly of kinetochores on centromeric chromatin to mediate interactions with spindle microtubules and control cell-cycle progression. MIS12 (minichromosome instability 12), also known as MTW1, hMis12, KNTC2AP or MIND kinetochore complex component homolog, is a 205 amino acid nuclear protein that is associated with the kinetochore. MIS12 is a component of the MIS12 complex, which is required for kinetochore formation during mitosis and normal chromosome alignment and segregation. The MIS12 complex consists of MIS12, DSN1, NSL1 and PMF-1. MIS12 is part of a network of complexes that provide microtubule attachment and generates pulling forces from depolymerization. MIS12 is encoded by a gene located on human chromosome 17, which comprises over 2.5% of the human genome and encodes over 1,200 genes.

## REFERENCES

- Goshima, G., Kiyomitsu, T., Yoda, K. and Yanagida, M. 2003. Human centromere chromatin protein hMIS12, essential for equal segregation, is independent of CENP-A loading pathway. *J. Cell Biol.* 160: 25-39.
- Obuse, C., Iwasaki, O., Kiyomitsu, T., Goshima, G., Toyoda, Y. and Yanagida, M. 2004. A conserved MIS12 centromere complex is linked to heterochromatic HP1 and outer kinetochore protein ZWINT-1. *Nat. Cell Biol.* 6: 1135-1141.
- Kline, S.L., Cheeseman, I.M., Hori, T., Fukagawa, T. and Desai, A. 2006. The human MIS12 complex is required for kinetochore assembly and proper chromosome segregation. *J. Cell Biol.* 173: 9-17.
- Liu, S.T., Rattner, J.B., Jablonski, S.A. and Yen, T.J. 2006. Mapping the assembly pathways that specify formation of the trilaminar kinetochore plates in human cells. *J. Cell Biol.* 175: 41-53.
- Zhang, R., Liu, S.T., Chen, W., Bonner, M., Pehrson, J., Yen, T.J. and Adams, P.D. 2007. HP1 proteins are essential for a dynamic nuclear response that rescues the function of perturbed heterochromatin in primary human cells. *Mol. Cell. Biol.* 27: 949-962.
- Hemmerich, P., Weidtkamp-Peters, S., Hoischen, C., Schmiedeberg, L., Erliandri, I. and Diekmann, S. 2008. Dynamics of inner kinetochore assembly and maintenance in living cells. *J. Cell Biol.* 180: 1101-1114.
- Wan, X., O'Quinn, R.P., Pierce, H.L., Joglekar, A.P., Gall, W.E., DeLuca, J.G., Carroll, C.W., Liu, S.T., Yen, T.J., McEwen, B.F., Stukenberg, P.T., Desai, A. and Salmon, E.D. 2009. Protein architecture of the human kinetochore microtubule attachment site. *Cell* 137: 672-684.
- Li, X. and Dawe, R.K. 2009. Fused sister kinetochores initiate the reductional division in meiosis I. *Nat. Cell Biol.* 11: 1103-1108.
- Joglekar, A.P., Bloom, K.S. and Salmon, E.D. 2010. Mechanisms of force generation by end-on kinetochore-microtubule attachments. *Curr. Opin. Cell Biol.* 22: 57-67.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## CHROMOSOMAL LOCATION

Genetic locus: MIS12 (human) mapping to 17p13.2; Mis12 (mouse) mapping to 11 B4.

## SOURCE

MIS12 (C-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of MIS12 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-107750 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

MIS12 (C-13) is recommended for detection of MIS12 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for MIS12 siRNA (h): sc-93889, MIS12 siRNA (m): sc-149443, MIS12 shRNA Plasmid (h): sc-93889-SH, MIS12 shRNA Plasmid (m): sc-149443-SH, MIS12 shRNA (h) Lentiviral Particles: sc-93889-V and MIS12 shRNA (m) Lentiviral Particles: sc-149443-V.

Molecular Weight of MIS12: 25 kDa.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.