

OAT3 (P-13): sc-107836

BACKGROUND

The organic anion transporter (OAT) family of proteins mediate the secretion of exogenous and endogenous metabolites from tissues throughout the body. OAT1 (organic anion transporter 1), a 563 amino acid protein, and OAT3 (organic anion transporter 3), a 542 amino acid protein, are 2 members of the OAT family and are highly expressed in kidneys. Localized specifically to the basolateral membrane, OAT1 and OAT3 are involved in the elimination of toxic organic anions, such as benzylpenicillin and cimetidine, from proximal renal tubules. Via their ability to remove anions from renal tissues, OAT1 and OAT3 are able to regulate the amount of toxins within the kidneys. Additionally, OAT1 functions as an organic anion exchanger that couples the uptake of one organic anion molecule with the efflux of one endogenous dicarboxylic acid molecule, such as ketoglutarate. Four isoforms of OAT1 and three isoforms of OAT3 are expressed due to alternative splicing events.

REFERENCES

- Lu, R., et al. 1999. Cloning of the human kidney PAH transporter: narrow substrate specificity and regulation by protein kinase C. *Am. J. Physiol.* 276: F295-F303.
- Race, J.E., et al. 1999. Molecular cloning and characterization of two novel human renal organic anion transporters (hOAT1 and hOAT3). *Biochem. Biophys. Res. Commun.* 255: 508-514.
- Bahn, A., et al. 2000. Genomic structure and *in vivo* expression of the human organic anion transporter 1 (hOAT1) gene. *Biochem. Biophys. Res. Commun.* 275: 623-630.
- Sun, W., et al. 2001. Isolation of a family of organic anion transporters from human liver and kidney. *Biochem. Biophys. Res. Commun.* 283: 417-422.
- Zhang, R., et al. 2008. Upregulation of rat renal cortical organic anion transporter (OAT1 and OAT3) expression in response to ischemia/reperfusion injury. *Am. J. Nephrol.* 28: 772-783.
- Vallon, V., et al. 2008. Overlapping *in vitro* and *in vivo* specificities of the organic anion transporters OAT1 and OAT3 for loop and thiazide diuretics. *Am. J. Physiol. Renal Physiol.* 294: F867-F873.
- Chen, J., et al. 2008. Adaptive responses of renal organic anion transporter 3 (OAT3) during cholestasis. *Am. J. Physiol. Renal Physiol.* 295: F247-F252.
- Hu, S., et al. 2008. Interaction of imatinib with human organic ion carriers. *Clin. Cancer Res.* 14: 3141-3148.
- Ogasawara, K., et al. 2008. Analysis of regulatory polymorphisms in organic ion transporter genes (SLC22A) in the kidney. *J. Hum. Genet.* 53: 607-614.

CHROMOSOMAL LOCATION

Genetic locus: Slc22a8 (mouse) mapping to 19 A.

SOURCE

OAT3 (P-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within a C-terminal extracellular domain of OAT3 of mouse origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-107836 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

OAT3 (P-13) is recommended for detection of OAT3 of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other OAT family members.

OAT3 (P-13) is also recommended for detection of OAT3 in additional species, including canine and porcine.

Suitable for use as control antibody for OAT3 siRNA (m): sc-150151, OAT3 shRNA Plasmid (m): sc-150151-SH and OAT3 shRNA (m) Lentiviral Particles: sc-150151-V.

Molecular Weight of OAT3: 62 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **OAT3 (3C11): sc-293264**, our highly recommended monoclonal alternative to OAT3 (P-13).