

▶ RBMS1 (C-14): sc-107995

BACKGROUND

RBMS1 (RNA binding motif, single stranded interacting protein 1), also known as YC1, MSSP (c-Myc single strand binding protein), SCR2 (suppressor of Cdc2 with RNA binding motif), MSSP-1, MSSP-2 or MSSP-3, is a member of the MSSP family of proteins. The MSSP family is comprised of proteins that bind to single stranded DNA/RNA. Through an interaction with the c-Myc protein, members of this family are involved in a wide variety of cellular functions, including gene transcription, DNA replication, apoptosis and cell cycle progression. RBMS1, a nuclear localized protein, is expressed in lung, placenta and heart with highest expression levels during the G₁ to S transition phase of the cell cycle. RBMS1 contains two RNP domains, namely RNP1-A and RNP1-B, both of which are necessary for DNA binding. RBMS1 specifically binds to a catalytic subunit of DNA polymerase (pol) α and stimulates its activity *in vitro*. Due to alternative splicing events, various isoforms exist for RBMS1.

REFERENCES

1. Kanaoka, Y. and Nojima, H. 1994. SCR: novel human suppressors of Cdc2/Cdc13 mutants of *Schizosaccharomyces pombe* harbour motifs for RNA binding proteins. *Nucleic Acids Res.* 22: 2687-2693.
2. Takai, T., et al. 1994. Molecular cloning of MSSP-2, a c-Myc gene single-strand binding protein: characterization of binding specificity and DNA replication activity. *Nucleic Acids Res.* 22: 5576-5581.
3. Negishi, Y., et al. 1994. Identification and cDNA cloning of single-stranded DNA binding proteins that interact with the region upstream of the human c-Myc gene. *Oncogene* 9: 1133-1143.
4. Haigermoser, C., et al. 1996. Cloning and characterization of the genomic DNA of the human MSSP genes. *Nucleic Acids Res.* 24: 3846-3857.
5. Niki, T., et al. 2000. MSSP, a protein binding to an origin of replication in the c-Myc gene, interacts with a catalytic subunit of DNA polymerase α and stimulates its polymerase activity. *FEBS Lett.* 475: 209-212.
6. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 602310. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
7. Fritz, D. and Stefanovic, B. 2007. RNA-binding protein RBMS3 is expressed in activated hepatic stellate cells and liver fibrosis and increases expression of transcription factor Prx1. *J. Mol. Biol.* 371: 585-595.

CHROMOSOMAL LOCATION

Genetic locus: RBMS1 (human) mapping to 2q24.2; Rbms1 (mouse) mapping to 2 C1.2.

SOURCE

RBMS1 (C-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of RBMS1 of human origin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-107995 X, 200 μ g/0.1 ml.

Blocking peptide available for competition studies, sc-107995 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

RBMS1 (C-14) is recommended for detection of RBMS1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with family members RBMS2 or RBMS3.

RBMS1 (C-14) is also recommended for detection of RBMS1 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for RBMS1 siRNA (h): sc-94943, RBMS1 siRNA (m): sc-152757, RBMS1 shRNA Plasmid (h): sc-94943-SH, RBMS1 shRNA Plasmid (m): sc-152757-SH, RBMS1 shRNA (h) Lentiviral Particles: sc-94943-V and RBMS1 shRNA (m) Lentiviral Particles: sc-152757-V.

RBMS1 (C-14) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Molecular Weight of RBMS1: 50 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfaction
Guaranteed

Try **RBMS1 (73-K2): sc-101190**, our highly recommended monoclonal alternative to RBMS1 (C-14).