

DAPK (H-300): sc-10805

BACKGROUND

DAPK (death associated protein) kinase and ZIP kinase are members of a novel protein kinase family, the members of which have the capacity to mediate apoptosis through their catalytic activities. DAP kinase contains a "death domain" and has been shown to mediate γ interferon-induced apoptosis. The introduction of DAP kinase into highly metastatic carcinoma clones lacking DAP kinase expression was shown to result in the suppression of metastasis, thus linking suppression of apoptosis to metastasis. ZIP kinase contains a leucine zipper domain, which is necessary for homodimerization and for interaction with other leucine zipper proteins. ZIP kinase dimerizes with ATF-4, an ATF/CREB transcription factor family member that contains a leucine zipper. Overexpression of ZIP kinase was shown to result in morphological changes associated with apoptosis in NIH/3T3 cells.

REFERENCES

1. Feinstein, E., et al. 1995. Assignment of DAP1 and DAPK—genes that positively mediate programmed cell death triggered by IFN- γ —to chromosome regions 5p12.2 and 9q34.1, respectively. *Genomics* 29: 305-307.
2. Sakagami, H., et al. 1997. Molecular cloning and developmental expression of a rat homologue of death-associated protein kinase in the nervous system. *Brain Res. Mol. Brain Res.* 52: 249-256.
3. Inbal, B., et al. 1997. DAP kinase links the control of apoptosis to metastasis. *Nature* 390: 180-184.
4. Kawai, T., et al. 1998. ZIP kinase, a novel serine/threonine kinase which mediates apoptosis. *Mol. Cell. Biol.* 18: 1642-1651.
5. Schumacher, A.M., et al. 2002. DAPK catalytic activity in the hippocampus increases during the recovery phase in an animal model of brain hypoxic-ischemic injury. *Biochim. Biophys. Acta* 1600: 128-137.
6. Jin, Y., et al. 2002. A death-associated protein kinase (DAPK)-interacting protein, DIP-1, is an E3 ubiquitin ligase that promotes tumor necrosis factor-induced apoptosis and regulates the cellular levels of DAPK. *J. Biol. Chem.* 277: 46980-46986.
7. Kim, W.S., et al. 2003. Promoter methylation and down-regulation of DAPK is associated with gastric atrophy. *Int. J. Mol. Med.* 12: 827-830.

CHROMOSOMAL LOCATION

Genetic locus: DAPK1 (human) mapping to 9q21.33.

SOURCE

DAPK (H-300) is a rabbit polyclonal antibody raised against amino acids 1132-1431 mapping at the C-terminus of DAP-kinase of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4°C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

DAPK (H-300) is recommended for detection of DAP-kinase of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000). DAPK (H-300) is also recommended for detection of DAP-kinase in additional species, including equine, canine, bovine, porcine and avian.


Suitable for use as control antibody for DAPK siRNA (h): sc-38976, DAPK shRNA Plasmid (h): sc-38976-SH and DAPK shRNA (h) Lentiviral Particles: sc-38976-V.

Molecular Weight of DAPK: 160 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2051 or ABC: sc-2018 rabbit IgG Staining Systems.

DATA



DAPK (H-300): sc-10805. Immunoperoxidase staining of formalin fixed, paraffin-embedded human duodenum tissue showing cytoplasmic and membrane staining of glandular cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **DAPK (17): sc-136286**, our highly recommended monoclonal alternative to DAPK (H-300).