

β-defensin 2 (M-17): sc-10858

BACKGROUND

β-defensins (also designated BD, and HBD in human) are small cationic peptides with broad-spectrum antimicrobial activity. β-defensins are involved in the resistance of epithelial surfaces, such as airway surface fluid, to microbial colonization. Human β-defensin 2 is locally regulated by inflammation and is the first member of the β-defensin family that is locally inducible by inflammation. The murine homolog of human β-defensin 2, which is called β-defensin 3, is present in the respiratory system and in low levels in the epithelial cells of the intestine and lung. The unique murine β-defensin 2 (Defβ2) is not expressed in airways of untreated mice, but is upregulated in the airways by lipopolysaccharide and may contribute to host defense at the mucosal surface of the airways.

REFERENCES

1. McCray, P.B., Jr., et al. 1997. Human airway epithelia express a β-defensin. *Am. J. Respir. Cell Mol. Biol.* 16: 343-349.
2. Liu, L., et al. 1997. The human β-defensin 1 and α-defensins are encoded by adjacent genes: two peptide families with differing disulfide topology share a common ancestry. *Genomics* 43: 316-320.
3. Liu, L., et al. 1998. Structure and mapping of the human β-defensin HBD-2 gene and its expression at sites of inflammation. *Gene* 222: 237-244.
4. Bals, R., et al. 1999. Mouse β-defensin 3 is an inducible antibacterial peptide expressed in the epithelia of multiple genes. *Infect. Immun.* 67: 3542-3547.
5. Yang, D., et al. 1999. β-defensins: linking innate and adaptive immunity through dendritic and T cell CCR-6. *Science* 286: 525-528.
6. Morrison, G.M., et al. 1999. A novel mouse β-defensin, Defβ2, which is upregulated in the airways by lipopolysaccharides. *FEBS Lett.* 442: 112-116.

CHROMOSOMAL LOCATION

Genetic locus: Defb2 (mouse) mapping to 8 A2.

SOURCE

β-defensin 2 (M-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of β-defensin 2 of mouse origin.

PRODUCT

Each vial contains 200 μg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-10858 P, (100 μg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

β-defensin 2 (M-17) is recommended for detection of precursor and mature β-defensin 2 of mouse origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for β-defensin 2 siRNA (m): sc-43722, β-defensin 2 shRNA Plasmid (m): sc-43722-SH and β-defensin 2 shRNA (m) Lentiviral Particles: sc-43722-V.

Molecular Weight of β-defensin 2: 5 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

SELECT PRODUCT CITATIONS

1. Selleri, S., et al. 2007. Toll-like receptor agonists regulate β-defensin 2 release in hair follicle. *Br. J. Dermatol.* 156: 1172-1177.
2. Gariboldi, S., et al. 2008. Low molecular weight hyaluronic acid increases the self-defense of skin epithelium by induction of β-defensin 2 via TLR2 and TLR4. *J. Immunol.* 181: 2103-2110.
3. Garreis, F., et al. 2010. Roles of human β-defensins in innate immune defense at the ocular surface: arming and alarming corneal and conjunctival epithelial cells. *Histochem. Cell Biol.* 134: 59-73.
4. Tohidnezhad, M., et al. 2011. Platelets display potent antimicrobial activity and release human β-defensin 2. *Platelets* 23: 217-223.
5. Mei, H.F., et al. 2012. β-defensin 2 as an adjuvant promotes anti-melanoma immune responses and inhibits the growth of implanted murine melanoma *in vivo*. *PLoS ONE* 7: e31328.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.