

Ribosomal Protein L13 (S-12): sc-109012

BACKGROUND

Ribosomes, the organelles that catalyze protein synthesis, are composed of a small subunit (40S) and a large subunit (60S) that consist of over 80 distinct ribosomal proteins. Mammalian ribosomal proteins are encoded by multigene families that contain processed pseudogenes and one functional intron-containing gene within their coding regions. Ribosomal Protein L13, also known as RPL13 or BBC1 (breast basic conserved protein 1), is a 211 amino acid protein that is a component of the 60S subunit. Localized to the cytoplasm and expressed ubiquitously, Ribosomal Protein L13 belongs to the L13E family of ribosomal proteins and functions in protein synthesis. In addition, the gene encoding Ribosomal Protein L13 is expressed at high levels in benign breast lesions. Like most ribosomal proteins, Ribosomal Protein L13 exists as multiple processed pseudogenes that are scattered throughout the genome. Due to alternative splicing events and/or alternative polyadenylation, various isoforms exist for Ribosomal Protein L13.

REFERENCES

- Adams, S.M., et al. 1992. Isolation and characterization of a novel gene with differential expression in benign and malignant human breast tumours. *Hum. Mol. Genet.* 1: 91-96.
- Bertauche, N., et al. 1994. Conservation of the human breast basic conserved 1 gene in the plant kingdom: characterization of a cDNA clone from *Arabidopsis thaliana*. *Gene* 141: 211-214.
- Moerland, E., et al. 1997. Exclusion of BBC1 and CMAR as candidate breast tumour-suppressor genes. *Br. J. Cancer* 76: 1550-1553.
- Kenmochi, N., et al. 1998. A map of 75 human ribosomal protein genes. *Genome Res.* 8: 509-523.
- Stubbs, A.P., et al. 1999. Differentially expressed genes in hormone refractory prostate cancer: association with chromosomal regions involved with genetic aberrations. *Am. J. Pathol.* 154: 1335-1343.
- Sáez-Vásquez, J., et al. 2000. Accumulation and nuclear targeting of BnC24, a *Brassica napus* ribosomal protein corresponding to a mRNA accumulating in response to cold treatment. *Plant Sci.* 156: 35-46.
- Jain, M., et al. 2004. Molecular characterization of a light-responsive gene, breast basic conserved protein 1 (OsiBBC1), encoding nuclear-localized protein homologous to Ribosomal Protein L13 from *Oryza sativa indica*. *Biochim. Biophys. Acta* 1676: 182-192.

CHROMOSOMAL LOCATION

Genetic locus: RPL13 (human) mapping to 16q24.3; Rpl13 (mouse) mapping to 8 E1.

SOURCE

Ribosomal Protein L13 (S-12) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Ribosomal Protein L13 of human origin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-109012 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Ribosomal Protein L13 (S-12) is recommended for detection of Ribosomal Protein L13 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Ribosomal Protein L13 (S-12) is also recommended for detection of Ribosomal Protein L13 in additional species, including equine, bovine and porcine.

Suitable for use as control antibody for Ribosomal Protein L13 siRNA (h): sc-92988, Ribosomal Protein L13 siRNA (m): sc-152894, Ribosomal Protein L13 shRNA Plasmid (h): sc-92988-SH, Ribosomal Protein L13 shRNA Plasmid (m): sc-152894-SH, Ribosomal Protein L13 shRNA (h) Lentiviral Particles: sc-92988-V and Ribosomal Protein L13 shRNA (m) Lentiviral Particles: sc-152894-V.

Molecular Weight of Ribosomal Protein L13: 24 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, K-562 whole cell lysate: sc-2203 or HeLa nuclear extract: sc-2120.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.


 MONOS
 Satisfaction
 Guaranteed

Try **Ribosomal Protein L13 (SS-09): sc-100829**, our highly recommended monoclonal alternative to Ribosomal Protein L13 (S-12).