

Syntaxin 8 (N-13): sc-109056

BACKGROUND

Syntaxins, a family of proteins involved in the fusion of synaptic vesicles with the plasma membrane, display broad tissue distribution and contain C-terminal hydrophobic domains that direct them to their respective intracellular compartments. Syntaxin 8, also known as STX8 or CARB, is a 236 amino acid single-pass type IV membrane protein that contains one t-SNARE coiled-coil homology domain and belongs to the Syntaxin family. Highly expressed in liver, brain, kidney, lung, placenta, spleen, pancreas and skeletal muscle, Syntaxin 8 functions in the early secretory pathway as a vesicle trafficking protein that shuttles proteins from the *cis*-Golgi membrane to the endoplasmic reticulum (ER). In addition, Syntaxin 8 associates with Syntaxin 7, v-SNARE Vti1p and endobrevin to form a SNARE complex that plays a role in the homo-typic fusion of late endosomes.

REFERENCES

1. Steegmaier, M., et al. 1998. Three novel proteins of the Syntaxin/SNAP 25 family. *J. Biol. Chem.* 273: 34171-34179.
2. Prekeris, R., et al. 1999. Differential roles of Syntaxin 7 and Syntaxin 8 in endosomal trafficking. *Mol. Biol. Cell* 10: 3891-3908.
3. Thoreau, V., et al. 1999. Molecular cloning, expression analysis, and chromosomal localization of human Syntaxin 8 (STX8). *Biochem. Biophys. Res. Commun.* 257: 577-583.
4. Online Mendelian Inheritance in Man, OMIM[™]. 1999. Johns Hopkins University, Baltimore, MD. MIM Number: 604203. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Bogdanovic, A., et al. 2002. Syntaxin 7, Syntaxin 8, Vti1 and VAMP-7 (vesicle-associated membrane protein 7) form an active SNARE complex for early macropinosytic compartment fusion in *Dictyostelium discoideum*. *Biochem. J.* 368: 29-39.

CHROMOSOMAL LOCATION

Genetic locus: STX8 (human) mapping to 17p13.1; Stx8 (mouse) mapping to 11 B3.

SOURCE

Syntaxin 8 (N-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an N-terminal cytoplasmic domain of Syntaxin 8 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-109056 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Syntaxin 8 (N-13) is recommended for detection of Syntaxin 8 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other Syntaxin family members.

Syntaxin 8 (N-13) is also recommended for detection of Syntaxin 8 in additional species, including equine, canine and bovine.

Suitable for use as control antibody for Syntaxin 8 siRNA (h): sc-93822, Syntaxin 8 siRNA (m): sc-153995, Syntaxin 8 shRNA Plasmid (h): sc-93822-SH, Syntaxin 8 shRNA Plasmid (m): sc-153995-SH, Syntaxin 8 shRNA (h) Lentiviral Particles: sc-93822-V and Syntaxin 8 shRNA (m) Lentiviral Particles: sc-153995-V.


Molecular Weight of Syntaxin 8: 27 kDa.

Positive Controls: A-431 whole cell lysate: sc-2201.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941. 3) Immunohistochemistry: use ImmunoCruz[™]: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

DATA



Syntaxin 8 (N-13): sc-109056. Immunoperoxidase staining of formalin fixed, paraffin-embedded human heart muscle tissue showing cytoplasmic staining of myocytes.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.