

# VAC14 (C-14): sc-109135

## BACKGROUND

Phosphatidylinositol 3,5-bisphosphate (PI(3,5)P<sub>2</sub>) is a signaling molecule that exists as a minor component of cell membranes and is essential for the distinguishing of cellular compartments. The synthesis of PI(3,5)P<sub>2</sub> is regulated by a number of proteins that are involved in intracellular trafficking and assembly events throughout the cell. VAC14, also known as TAX1BP2 (Tax1-binding protein 2) or TRX, is a 782 amino acid protein that contains 6 HEAT repeats and exists as part of a regulatory complex with FIG4. Expressed ubiquitously, VAC14 works with FIG4 to control the synthesis of PI(3,5)P<sub>2</sub>, specifically mediating the activation of PIP5KIII, a kinase involved in the regulation of PI(3,5)P<sub>2</sub> activity. The gene encoding VAC14 maps to human chromosome 16, which houses over 900 genes and comprises nearly 3% of the human genome.

## REFERENCES

- Mireskandari, A., et al. 1996. Isolation of a cDNA clone, TRX encoding a human T cell lymphotropic virus type-I Tax1 binding protein. *Biochim. Biophys. Acta* 1306: 9-13.
- Sbrissa, D., et al. 2004. A mammalian ortholog of *Saccharomyces cerevisiae* Vac14 that associates with and upregulates PIKfyve phosphoinositide 5-kinase activity. *Mol. Cell. Biol.* 24: 10437-10447.
- Lemaire, J.F. and McPherson, P.S. 2006. Binding of VAC14 to neuronal nitric oxide synthase: Characterisation of a new internal PDZ-recognition motif. *FEBS Lett.* 580: 6948-6954.
- Ching, Y.P., et al. 2006. The retroviral oncoprotein Tax targets the coiled-coil centrosomal protein TAX1BP2 to induce centrosome overduplication. *Nat. Cell Biol.* 8: 717-724.
- Sbrissa, D., et al. 2007. Core protein machinery for mammalian phosphatidylinositol 3,5-bisphosphate synthesis and turnover that regulates the progression of endosomal transport. Novel Sac phosphatase joins the ArPIKfyve-PIKfyve complex. *J. Biol. Chem.* 282: 23878-23891.
- Zhang, Y., et al. 2007. Loss of VAC14, a regulator of the signaling lipid phosphatidylinositol 3,5-bisphosphate, results in neurodegeneration in mice. *Proc. Natl. Acad. Sci. USA* 104: 17518-17523.
- Jin, N., et al. 2008. VAC14 nucleates a protein complex essential for the acute interconversion of PI3P and PI(3,5)P<sub>2</sub> in yeast and mouse. *EMBO J.* 27: 3221-3234.
- Online Mendelian Inheritance in Man, OMIM™. 2008. Johns Hopkins University, Baltimore, MD. MIM Number: 604632. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

## CHROMOSOMAL LOCATION

Genetic locus: VAC14 (human) mapping to 16q22.1; Vac14 (mouse) mapping to 8 E1.

## SOURCE

VAC14 (C-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of VAC14 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-109135 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

VAC14 (C-14) is recommended for detection of VAC14 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

VAC14 (C-14) is also recommended for detection of VAC14 in additional species, including equine, canine and bovine.

Suitable for use as control antibody for VAC14 siRNA (h): sc-72206, VAC14 siRNA (m): sc-72207, VAC14 shRNA Plasmid (h): sc-72206-SH, VAC14 shRNA Plasmid (m): sc-72207-SH, VAC14 shRNA (h) Lentiviral Particles: sc-72206-V and VAC14 shRNA (m) Lentiviral Particles: sc-72207-V.

Molecular Weight of VAC14: 88 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, K-562 whole cell lysate: sc-2203 or Hep G2 cell lysate: sc-2227.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.