

ZNF207 (P-14): sc-109180



The Power to Question

BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. ZNF207 (zinc-finger protein 207) is a 478 amino acid protein that localizes to the nucleus and contains 2 C₂H₂-type zinc fingers. Expressed ubiquitously, ZNF207 may function as a transcription factor. Three isoforms of ZNF207 are expressed due to alternative splicing events.

REFERENCES

1. Rousseau-Merck, M.F., et al. 1994. Chromosomal localization of 9 KOX zinc-finger genes: physical linkages suggest clustering of KOX genes on chromosomes 12, 16, and 19. *Hum. Genet.* 92: 583-587.
2. Pahl, P.M., et al. 1998. ZNF207, a ubiquitously expressed zinc-finger gene on chromosome 6p21.3. *Genomics* 53: 410-412.
3. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 603428. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Sun, Y., et al. 2003. The KRAB domain of zinc-finger gene ZNF268: a potential transcriptional repressor. *IUBMB Life* 55: 127-131.
5. Rousseau-Merck, M.F., et al. 2003. The KOX zinc-finger genes: genome wide mapping of 368 ZNF PAC clones with zinc-finger gene clusters predominantly in 23 chromosomal loci are confirmed by human sequences annotated in EnsEMBL. *Cytogenet. Genome Res.* 98: 147-153.
6. Nakamura, M., et al. 2004. A novel subfamily of zinc-finger genes involved in embryonic development. *J. Cell. Biochem.* 93: 887-895.
7. Englbrecht, C.C., et al. 2004. Conservation, diversification and expansion of C₂H₂ zinc finger proteins in the *Arabidopsis thaliana* genome. *BMC Genomics* 5: 39.
8. O'Geen, H., et al. 2007. Genome-wide analysis of KAP1 binding suggests autoregulation of KRAB-ZNFs. *PLoS Genet.* 3: e89.

CHROMOSOMAL LOCATION

Genetic locus: ZNF207 (human) mapping to 17q11.2; Zfp207 (mouse) mapping to 11 B5.

SOURCE

ZNF207 (P-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of ZNF207 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-109180 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

ZNF207 (P-14) is recommended for detection of ZNF207 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other ZNF family members.

ZNF207 (P-14) is also recommended for detection of ZNF207 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for ZNF207 siRNA (h): sc-93847, ZNF207 siRNA (m): sc-155654, ZNF207 shRNA Plasmid (h): sc-93847-SH, ZNF207 shRNA Plasmid (m): sc-155654-SH, ZNF207 shRNA (h) Lentiviral Particles: sc-93847-V and ZNF207 shRNA (m) Lentiviral Particles: sc-155654-V.


Molecular Weight of ZNF207: 51 kDa.

Positive Controls: Hep G2 nuclear extract: sc-364819, HeLa nuclear extract: sc-2120 or Hep G2 cell lysate: sc-2227.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



ZNF207 (P-14): sc-109180. Western blot analysis of ZNF207 expression in Hep G2 nuclear extract.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.