

ASPP1 (M-18): sc-10921

BACKGROUND

ASPP proteins interact with p53 and are responsible for enhancing p53-induced apoptosis but not cell cycle arrest. Inhibition of endogenous ASPP1 (PPP1R13B) function inhibits the apoptotic function of endogenous p53 in response to apoptotic stimuli. ASPP1 amplifies DNA binding and transactivation function of p53 on the promoters of proapoptotic genes *in vivo*. Expression of ASPP1 is often downregulated in human breast carcinomas expressing wildtype p53, but not in those expressing mutant p53. This research indicates that ASPP1 regulates the tumor suppression function of p53 *in vivo*. ASPP1 is predominantly a cytoplasmic protein, although some fraction of the polypeptide is nuclear. Defects in PPP1R13B, the gene which encodes ASPP1, may be a cause of breast cancers. The deduced ASPP1 protein contains 1,090 amino acid residues.

REFERENCES

1. Nagase, T., et al. 1999. Prediction of the coding sequences of unidentified human genes. XI. The complete sequences of 100 new cDNA clones from brain which code for large proteins *in vitro*. DNA Res. 5: 277-286.
2. Samuels-Lev, Y., et al. 2001. ASPP proteins specifically stimulate the apoptotic function of p53. Mol. Cell 8: 781-794.
3. Bergamaschi, D., et al. 2004. ASPP1 and ASPP2: common activators of p53 family members. Mol. Cell. Biol. 24: 1341-1350.
4. Bergamaschi, D., et al. 2005. MDM2 and MDMX prevent ASPP1 and ASPP2 from stimulating p53 without targeting p53 for degradation. Oncogene 24: 3836-3841.
5. Fogal, V., et al. 2005. ASPP1 and ASPP2 are new transcriptional targets of E2F. Cell Death Differ. 12: 369-376.

CHROMOSOMAL LOCATION

Genetic locus: PPP1R13B (human) mapping to 14q32.33; Ppp1r13b (mouse) mapping to 12 F1.

SOURCE

ASPP1 (M-18) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of ASPP1 of mouse origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-10921 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

ASPP1 (M-18) is recommended for detection of ASPP1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


ASPP1 (M-18) is also recommended for detection of ASPP1 in additional species, including equine, canine and avian.

Suitable for use as control antibody for ASPP1 siRNA (h): sc-60214, ASPP1 siRNA (m): sc-60215, ASPP1 shRNA Plasmid (h): sc-60214-SH, ASPP1 shRNA Plasmid (m): sc-60215-SH, ASPP1 shRNA (h) Lentiviral Particles: sc-60214-V and ASPP1 shRNA (m) Lentiviral Particles: sc-60215-V.

Molecular Weight of ASPP1: 119 kDa.

Positive Control: Saos-2 cell lysate: sc-2235, HeLa whole cell lysate: sc-2200 or WI-38 whole cell lysate: sc-364260.

DATA



ASPP1 (M-18): sc-10921. Western blot analysis of ASPP1 expression in HeLa (A) and WI-38 (B) whole cell lysates.

SELECT PRODUCT CITATIONS

1. Liu, F., et al. 2007. Insight into the host-parasite interplay by proteomic study of host proteins copurified with the human parasite, *Schistosoma japonicum*. Proteomics 7: 450-462.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **ASPP1 (LX011): sc-53903**, our highly recommended monoclonal alternative to ASPP1 (M-18).