



MBPH (N-17): sc-109320

BACKGROUND

MBPH, also known as PRG3 (proteoglycan 3), is a 225 amino acid protein that contains one C-type lectin domain and localizes to cytoplasmic granules. Expressed in bone marrow, MBPH functions to stimulate neutrophil superoxide production and IL-8 release, possibly playing a role in cytotoxic and cytostimulatory activities within the cell. The gene encoding MBPH maps to human chromosome 11, which comprises approximately 4% of human genomic DNA and is considered a gene and disease association dense chromosome. The chromosome 11 encoded *Atm* gene is important for regulation of cell cycle arrest and apoptosis following double strand DNA breaks. *Atm* mutation leads to the disorder known as ataxia-telangiectasia. The blood disorders Sickle cell anemia and thalassemia are caused by *HBB* gene mutations, while Wilms' tumors, WAGR syndrome and Denys-Drash syndrome are associated with mutations of the *WT1* gene. Jervell and Lange-Nielsen syndrome, Jacobsen syndrome, Niemann-Pick disease, hereditary angioedema and Smith-Lemli-Opitz syndrome are also associated with defects in chromosome 11-encoded genes.

REFERENCES

- Plager, D.A., et al. 1999. A novel and highly divergent homolog of human eosinophil granule major basic protein. *J. Biol. Chem.* 274: 14464-14473.
- Yerushalmi, G.M., et al. 2002. Met-HGF/SF signal transduction induces mimp, a novel mitochondrial carrier homologue, which leads to mitochondrial depolarization. *Neoplasia.* 4: 510-522.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 606814. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Gross, A. 2005. Mitochondrial carrier homolog 2: a clue to cracking the BCL-2 family riddle? *J. Bioenerg. Biomembr.* 37: 113-119.
- Grinberg, M., et al. 2005. Mitochondrial carrier homolog 2 is a target of tBID in cells signaled to die by tumor necrosis factor alpha. *Mol. Cell. Biol.* 25: 4579-4590.
- Schwarz, M., et al. 2007. Mitochondrial carriers and pores: key regulators of the mitochondrial apoptotic program? *Apoptosis.* 12: 869-876.
- Ataga, K.I., et al. 2007. β -thalassaemia and sickle cell anaemia as paradigms of hypercoagulability. *Br. J. Haematol.* 139: 3-13.
- Berger, A.C., et al. 2007. The subcellular localization of the Niemann-Pick Type C proteins depends on the adaptor complex AP-3. *J. Cell. Sci.* 120: 3640-3652.
- Hanein, S., et al. 2009. TMEM126A, encoding a mitochondrial protein, is mutated in autosomal-recessive nonsyndromic optic atrophy. *Am. J. Hum. Genet.* 84: 493-498.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

CHROMOSOMAL LOCATION

Genetic locus: PRG3 (human) mapping to 11q12.1.

SOURCE

MBPH (N-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of MBPH of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-109320 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

MBPH (N-17) is recommended for detection of MBPH of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for MBPH siRNA (h): sc-96304, MBPH shRNA Plasmid (h): sc-96304-SH and MBPH shRNA (h) Lentiviral Particles: sc-96304-V.

Molecular Weight of pre-pro form MBPH: 25 kDa.

Molecular Weight of active peptide MBPH: 14 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.