

PCDHB3 (T-12): sc-109777

BACKGROUND

Protocadherins are a large family of cadherin-like cell adhesion proteins that are involved in the establishment and maintenance of neuronal connections in the brain. There are three protocadherin gene clusters, designated α , β and γ , all of which contain multiple tandemly arranged genes. PCDHB3 (protocadherin β -3) is a 796 amino acid single pass transmembrane protein that is one of 16 proteins in the protocadherin beta cluster. Unlike the alpha and γ gene clusters whose genes are spliced to downstream constant region exons during transcription, members of the β cluster (such as PCDHB3) do not use constant-region exons to produce mRNAs. As a result, each protocadherin β gene encodes the transmembrane, extracellular and short cytoplasmic domains of the protein. PCDHB3 is likely a calcium-dependent cell adhesion protein that is involved in the maintenance of neural connections in the brain.

REFERENCES

1. Wu, Q. and Maniatis, T. 1999. A striking organization of a large family of human neural cadherin-like cell adhesion genes. *Cell* 97: 779-790.
2. Yagi, T. and Takeichi, M. 2000. Cadherin superfamily genes: functions, genomic organization, and neurologic diversity. *Genes Dev.* 14: 1169-1180.
3. Vanhalst, K., Kools, P., Vanden Eynde, E. and van Roy, F. 2001. The human and murine protocadherin- β one-exon gene families show high evolutionary conservation, despite the difference in gene number. *FEBS Lett.* 495: 120-125.
4. Wu, Q., Zhang, T., Cheng, J.F., Kim, Y., Grimwood, J., Schmutz, J., Dickson, M., Noonan, J.P., Zhang, M.Q., Myers, R.M. and Maniatis, T. 2001. Comparative DNA sequence analysis of mouse and human protocadherin gene clusters. *Genome Res.* 11: 389-404.
5. Hamada, S. and Yagi, T. 2001. The cadherin-related neuronal receptor family: a novel diversified cadherin family at the synapse. *Neurosci. Res.* 41: 207-215.
6. Morishita, H. and Yagi, T. 2007. Protocadherin family: diversity, structure, and function. *Curr. Opin. Cell Biol.* 19: 584-592.
7. Online Mendelian Inheritance in Man, OMIM[™]. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 606329. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
8. Yagi, T. 2008. Clustered protocadherin family. *Dev. Growth Differ.* 50: S131-S140.

CHROMOSOMAL LOCATION

Genetic locus: PCDHB3 (human) mapping to 5q31.3; Pcdhb3 (mouse) mapping to 18 B3.

SOURCE

PCDHB3 (T-12) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an extracellular domain of PCDHB3 of human origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-109777 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

PCDHB3 (T-12) is recommended for detection of PCDHB3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other PCDHB family members.

PCDHB3 (T-12) is also recommended for detection of PCDHB3 in additional species, including equine.

Suitable for use as control antibody for PCDHB3 siRNA (h): sc-91946, Pcdhb3 siRNA (m): sc-152078, PCDHB3 shRNA Plasmid (h): sc-91946-SH, Pcdhb3 shRNA Plasmid (m): sc-152078-SH, PCDHB3 shRNA (h) Lentiviral Particles: sc-91946-V and Pcdhb3 shRNA (m) Lentiviral Particles: sc-152078-V.

Molecular Weight of PCDHB3: 87 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.