## SANTA CRUZ BIOTECHNOLOGY, INC.

# PCDHGA11 (G-13): sc-109806



#### BACKGROUND

Protocadherins are a large family of cadherin-like cell adhesion proteins that are involved in the establishment and maintenance of neuronal connections in the brain. There are three protocadherin (PCDH) gene clusters, designated  $\alpha$ ,  $\beta$  and  $\gamma$ , all of which contain multiple tandemly arranged genes. PCD-HGA11 (protocadherin  $\gamma$ -A11) is a 935 amino acid that is one of 22 proteins encoded by the protocadherin  $\gamma$  cluster. The protocadherein  $\gamma$  cluster consists of three subfamilies (A, B and C) and PCDHGA11 is a member of the  $\gamma$  subfamily A. PCDHGA11 is a type I transmembrane receptor containing six cadherin motifs and is expressed in the central nervous system where it localizes to synapses. Members of the  $\gamma$  cluster of protocadherins are essential for neuronal survival. There are two isoforms of PCDHGA11 that are produced as a result of alternative splicing events.

## REFERENCES

- 1. Kohmura, N., et al. 1998. Diversity revealed by a novel family of cadherins expressed in neurons at a synaptic complex. Neuron 20: 1137-1151.
- 2. Wu, Q., et al. 2001. Comparative DNA sequence analysis of mouse and human protocadherin gene clusters. Genome Res. 11: 389-404.
- 3. Tasic, B., et al. 2002. Promoter choice determines splice site selection in protocadherin  $\alpha$  and  $\gamma$  pre-mRNA splicing. Mol. Cell 10: 21-33.
- Wang, X., et al. 2002. γ protocadherins are required for survival of spinal interneurons. Neuron 36: 843-854.
- 5. Kirov, G., et al. 2003. Variation in the protocadherin  $\gamma$  A gene cluster. Genomics 82: 433-440.
- Zou, C., et al. 2007. Sequence analysis and expression mapping of the rat clustered protocadherin gene repertoires. Neuroscience 144: 579-603.
- 7. Online Mendelian Inheritance in Man, OMIM™. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 606297. World Wide Web URL: http://www.ncbi.nlm.nih.gov/omim/
- Dallosso, A.R., et al. 2009. Frequent long-range epigenetic silencing of protocadherin gene clusters on chromosome 5q31 in Wilms' tumor. PLoS Genet. 5: e1000745.

## CHROMOSOMAL LOCATION

Genetic locus: PCDHGA11 (human) mapping to 5q31.3.

## SOURCE

PCDHGA11 (G-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an extracellular domain of PCDHGA11 of human origin.

### PRODUCT

Each vial contains 200  $\mu g$  IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-109806 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

### APPLICATIONS

PCDHGA11 (G-13) is recommended for detection of PCDHGA11 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other PCDHGA family members.

PCDHGA11 (G-13) is also recommended for detection of PCDHGA11 in additional species, including canine.

Suitable for use as control antibody for PCDHGA11 siRNA (h): sc-91755, PCDHGA11 shRNA Plasmid (h): sc-91755-SH and PCDHGA11 shRNA (h) Lentiviral Particles: sc-91755-V.

Molecular Weight of PCDHGA11: 102 kDa.

### **RECOMMENDED SECONDARY REAGENTS**

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluo-rescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

#### **STORAGE**

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

#### PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.