

# PCDHGA9 (V-13): sc-109839

## BACKGROUND

Protocadherins are a large family of cadherin-like cell adhesion proteins that are involved in the establishment and maintenance of neuronal connections in the brain. There are three protocadherin (PCDH) gene clusters, designated  $\alpha$ ,  $\beta$  and  $\gamma$ , all of which contain multiple tandemly arranged genes. PCDHGA9 (protocadherin  $\gamma$ -A9) is a 932 amino acid that is one of 22 proteins encoded by the protocadherin  $\gamma$  cluster. The protocadherin  $\gamma$  cluster consists of three subfamilies (A, B and C) and PCDHGA9 is a member of the  $\gamma$  subfamily A. PCDHGA9 is a type I transmembrane receptor containing six cadherin motifs and is expressed in the central nervous system where it localizes to synapses. Members of the  $\gamma$  cluster of protocadherins are essential for neuronal survival. There are two isoforms of PCDHGA9 that are produced as a result of alternative splicing events.

## REFERENCES

1. Kohmura, N., et al. 1998. Diversity revealed by a novel family of cadherins expressed in neurons at a synaptic complex. *Neuron* 20: 1137-1151.
2. Wu, Q., et al. 2001. Comparative DNA sequence analysis of mouse and human protocadherin gene clusters. *Genome Res.* 11: 389-404.
3. Tasic, B., et al. 2002. Promoter choice determines splice site selection in protocadherin  $\alpha$  and  $\gamma$  pre-mRNA splicing. *Mol. Cell.* 10: 21-33.
4. Wang, X., et al. 2002.  $\gamma$  protocadherins are required for survival of spinal interneurons. *Neuron* 36: 843-854.
5. Kirov, G., et al. 2003. Variation in the protocadherin  $\gamma$  A gene cluster. *Genomics* 82: 433-440.
6. Zou, C., et al. 2007. Sequence analysis and expression mapping of the rat clustered protocadherin gene repertoires. *Neuroscience* 144: 579-603.
7. Online Mendelian Inheritance in Man, OMIM™. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 606297. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
8. Dallosso, A.R., et al. 2009. Frequent long-range epigenetic silencing of protocadherin gene clusters on chromosome 5q31 in Wilms' tumor. *PLoS Genet.* 5: e1000745.

## CHROMOSOMAL LOCATION

Genetic locus: Pcdhga9 (mouse) mapping to 18 B3; Pcdhga9 (rat) mapping to 18p11.

## SOURCE

PCDHGA9 (V-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of PCDHGA9 of mouse origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-109839 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

PCDHGA9 (V-13) is recommended for detection of PCDHGA9 of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other PCDHGA family members.

Suitable for use as control antibody for Pcdhga9 siRNA (m): sc-152096, Pcdhga9 shRNA Plasmid (m): sc-152096-SH and Pcdhga9 shRNA (m) Lentiviral Particles: sc-152096-V.

Molecular Weight of PCDHGA9 isoforms: 102/91 kDa.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.