

HS6ST1 (C-14): sc-109941

BACKGROUND

Heparan sulfate structures, which are responsible for executing multiple biologic activities, are generated and regulated by heparan sulfate (HS) biosynthetic enzymes. HS6ST1 (heparan sulfate 6-O-sulfotransferase 1), also known as HS6ST, is a 411 amino acid single-pass type II membrane protein that exists as multiple alternatively spliced isoforms and belongs to the sulfotransferase 6 family. Expressed in fetal brain, HS6ST1 functions as a 6-O-sulfation enzyme that specifically catalyzes the transfer of sulfate from 3'-phosphoadenosine 5'-phosphosulfate (PAPS) to the N-sulfoglucosamine residue (GlcNS) HS. HS6ST1 is subject to post-translational N-glycosylation and is encoded by a functional gene on human chromosome 2 and a pseudogene on human chromosome 1 (known as LOC728969).

REFERENCES

- Habuchi, H., Kobayashi, M. and Kimata, K. 1998. Molecular characterization and expression of heparan-sulfate 6-sulfotransferase. Complete cDNA cloning in human and partial cloning in Chinese hamster ovary cells. *J. Biol. Chem.* 273: 9208-9213.
- Habuchi, H., Tanaka, M., Habuchi, O., Yoshida, K., Suzuki, H., Ban, K. and Kimata, K. 2000. The occurrence of three isoforms of heparan sulfate 6-O-sulfotransferase having different specificities for hexuronic acid adjacent to the targeted N-sulfoglucosamine. *J. Biol. Chem.* 275: 2859-2868.
- Habuchi, H., Miyake, G., Nogami, K., Kuroiwa, A., Matsuda, Y., Kusche-Gullberg, M., Habuchi, O., Tanaka, M. and Kimata, K. 2003. Biosynthesis of heparan sulphate with diverse structures and functions: two alternatively spliced forms of human heparan sulphate 6-O-sulphotransferase-2 having different expression patterns and properties. *Biochem. J.* 371: 131-142.
- Edavettal, S.C., Carrick, K., Shah, R.R., Pedersen, L.C., Tropsha, A., Pope, R.M. and Liu, J. 2004. A conformational change in heparan sulfate 3-O-sulfotransferase-1 is induced by binding to heparan sulfate. *Biochemistry* 43: 4680-4688.
- Online Mendelian Inheritance in Man, OMIM™. 2005. Johns Hopkins University, Baltimore, MD. MIM Number: 604846. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Backen, A.C., Cole, C.L., Lau, S.C., Clamp, A.R., McVey, R., Gallagher, J.T. and Jayson, G.C. 2007. Heparan sulphate synthetic and editing enzymes in ovarian cancer. *Br. J. Cancer* 96: 1544-1548.
- Nagai, N., Habuchi, H., Kitazume, S., Toyoda, H., Hashimoto, Y. and Kimata, K. 2007. Regulation of heparan sulfate 6-O-sulfation by β -secretase activity. *J. Biol. Chem.* 282: 14942-14951.

CHROMOSOMAL LOCATION

Genetic locus: HS6ST1 (human) mapping to 2q14.3; Hs6st1 (mouse) mapping to 1 B.

SOURCE

HS6ST1 (C-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of HS6ST1 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-109941 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

HS6ST1 (C-14) is recommended for detection of HS6ST1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with family members HS6ST2 or HS6ST3.

HS6ST1 (C-14) is also recommended for detection of HS6ST1 in additional species, including equine.

Suitable for use as control antibody for HS6ST1 siRNA (h): sc-94375, HS6ST1 siRNA (m): sc-146089, HS6ST1 shRNA Plasmid (h): sc-94375-SH, HS6ST1 shRNA Plasmid (m): sc-146089-SH, HS6ST1 shRNA (h) Lentiviral Particles: sc-94375-V and HS6ST1 shRNA (m) Lentiviral Particles: sc-146089-V.

Molecular Weight (predicted) of HS6ST1: 48 kDa.

Molecular Weight (observed) of HS6ST1: 43 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **HS6ST1 (C-9): sc-398231**, our highly recommended monoclonal alternative to HS6ST1 (C-14).