

LMO4 (S-17): sc-11120

BACKGROUND

The LIM-only (LMO) proteins, LMO1 and LMO2, are nuclear factors that are characterized by a conserved LIM domain. The LIM domain consists of a cysteine-rich zinc-binding motif that is present in a variety of transcription factors, including the LIM homeobox (LHX) proteins expressed in the central nervous system and involved in cell differentiation. LMO1 and LMO2 are expressed in the adult CNS in a cell type-specific manner, where they are differentially regulated by neuronal activity and are involved in regulating the cellular differentiated phenotype of neurons. LMO2 lacks a specific DNA-binding homeobox domain but rather assembles into transcriptional regulatory complexes to mediate gene expression by interacting with the widely expressed nuclear LIM interactor (NLI). NLI, known also as CLIM-1, and the related protein CLIM-2 facilitate the formation of heteromeric LIM complexes and also enhance the nuclear retention of LIM proteins. LMO2 and the related protein LMO4 are expressed in thymic precursor cells. LMO4 is also expressed in mature T cells, cranial neural crest cells, somite, dorsal limb bud mesenchyme, motor neurons, and Schwann cell progenitors.

REFERENCES

- Boehm, T., et al. 1991. The rhombotin family of cysteine-rich LIM-domain oncogenes: distinct members are involved in T-cell translocations to human chromosomes 11p15 and 11p13. *Proc. Natl. Acad. Sci. USA* 88: 4367-4371.
- Agulnick, A.D., et al. 1996. Interactions of the LIM-domain-binding factor Ldb1 with LIM homeodomain proteins. *Nature* 384: 270-272.
- Hinks, G.L., et al. 1997. Expression of LIM protein genes LMO1, LMO2, and LMO3 in adult mouse hippocampus and other forebrain regions: differential regulation by seizure activity. *J. Neurosci.* 17: 5549-5559.
- Rabbitts, T.H., et al. 1997. Chromosomal translocations and leukaemia: a role for LMO2 in T cell acute leukaemia, in transcription and in erythropoiesis. *Leukemia* 3: 271-272.

CHROMOSOMAL LOCATION

Genetic locus: LMO4 (human) mapping to 1p22.3; Lmo4 (mouse) mapping to 3 H2.

SOURCE

LMO4 (C-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of LMO4 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-11120 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

LMO4 (S-17) is recommended for detection of LMO4 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

LMO4 (S-17) is also recommended for detection of LMO4 in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for LMO4 siRNA (h): sc-38029, LMO4 siRNA (m): sc-38030, LMO4 shRNA Plasmid (h): sc-38029-SH, LMO4 shRNA Plasmid (m): sc-38030-SH, LMO4 shRNA (h) Lentiviral Particles: sc-38029-V and LMO4 shRNA (m) Lentiviral Particles: sc-38030-V.

Molecular Weight of LMO4: 17 kDa.

Positive Controls: IMR-32 cell lysate: sc-2409.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

SELECT PRODUCT CITATIONS

- Leuba, G., et al. 2004. Differential expression of LMO4 protein in Alzheimer's disease. *Neuropathol. Appl. Neurobiol.* 30: 57-69.
- Leuba, G., et al. 2008. Postsynaptic density protein PSD-95 expression in Alzheimer's disease and okadaic acid induced neuritic retraction. *Neurobiol. Dis.* 30: 408-419.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **LMO4 (4H8): sc-293440**, our highly recommended monoclonal alternative to LMO4 (S-17).