

## PTPβ (C-20): sc-1114

### BACKGROUND

Protein tyrosine phosphatases, or PTPs, are type I transmembrane proteins, membrane associated proteins or proteins localized in nuclei. Examples of transmembrane PTPs are LAR, PTPα, PTPβ, PTPγ, PTPδ, PTPε, PTPζ, PTPκ and PTPμ. Transmembrane PTPs play diverse roles during development and in adult tissues. Immunodepletion studies have suggested LAR to be a regulator of Insulin receptor phosphorylation. PTPα activity is increased twofold in response to phorbol ester stimulation, resulting in serine phosphorylation either directly or indirectly by members of the PKC family. Overexpression of v-H-ras and Neu, but not Myc or Int2, in mammary tumors has been shown to induce PTPε expression. An alternative splicing event leads to a nervous tissue-specific chondroitin sulfate proteoglycan called phosphacan, which represents the amino terminal portion of PTPζ. PTPκ and PTPμ share a conserved amino terminal 160 amino acid MAM domain which facilitates homophilic binding. PTPμ localizes to points of cell contact and may be involved in regulating the assembly and disassembly of cadherin/catenin complexes *in vivo*.

### REFERENCES

- Ahmad, F., et al. 1995. Increased abundance of the receptor-type protein-tyrosine phosphatase LAR accounts for the elevated Insulin receptor dephosphorylating activity in adipose tissue of obese human subjects. *J. Clin. Invest.* 95: 2806-2812.
- den Hertog, J., et al. 1995. Stimulation of receptor protein-tyrosine phosphatase α activity and phosphorylation by phorbol ester. *Cell Growth Differ.* 6: 303-307.
- Brady-Kalnay, S.M., et al. 1995. Receptor protein tyrosine phosphatase PTPm associates with cadherins and catenins *in vivo*. *J. Cell. Biol.* 130: 977-986.
- Zondag, G.C., et al. 1995. Homophilic interactions mediated by receptor tyrosine phosphatases μ and κ. A critical role for the novel extracellular MAM domain. *J. Biol. Chem.* 270: 14247-14250.
- Milev, P., et al. 1995. Complex-type asparagine-linked oligosaccharides on phosphacan and protein-tyrosine phosphatase-ζ/β mediate their binding to neural cell adhesion molecules and tenascin. *J. Biol. Chem.* 270: 24650-24653.
- Elson, A., et al. 1995. Protein-tyrosine phosphatase ε. An isoform specifically expressed in mouse mammary tumors initiated by v-Ha-Ras or Neu. *J. Biol. Chem.* 270: 26116-26122.

### CHROMOSOMAL LOCATION

Genetic locus: PTPRB (human) mapping to 12q15.

### SOURCE

PTPβ (C-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of PTPβ of human origin.

### STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### PRODUCT

Each vial contains 200 μg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-1114 P, (100 μg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

### APPLICATIONS

PTPβ (C-20) is recommended for detection of PTPβ of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μg per 100-500 μg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


PTPβ (C-20) is also recommended for detection of PTPβ in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for PTPβ siRNA (h): sc-44051, PTPβ shRNA Plasmid (h): sc-44051-SH and PTPβ shRNA (h) Lentiviral Particles: sc-44051-V.

Molecular Weight of PTPβ: 200 kDa.

Positive Controls: A-431 whole cell lysate: sc-2201.

### DATA



PTPβ (C-20): sc-1114. Western blot analysis of processed PTPβ expression in A-431 whole cell lysate.

### SELECT PRODUCT CITATIONS

- Wu, C.W., et al. 2006. Protein tyrosine-phosphatase expression profiling in gastric cancer tissues. *Cancer Lett.* 242: 95-103.
- González-Fernández, L., et al. 2009. Identification of protein tyrosine phosphatases and dual-specificity phosphatases in mammalian spermatozoa and their role in sperm motility and protein tyrosine phosphorylation. *Biol. Reprod.* 80: 1239-1252.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.