

Topo III β -1,2,3 (D-16): sc-11264

BACKGROUND

DNA topoisomerases are nuclear enzymes that regulate the topological structure of DNA by transiently breaking and rejoining DNA strands. Although DNA topoisomerase I and DNA topoisomerase II relax both positive and negative supercoils, DNA topoisomerase III relaxes only negative supercoils. DNA topoisomerase III α exists as a long and a short isoform, which are produced by alternative splicing. DNA topoisomerase III α , which localizes to the nucleolus, is constitutively expressed and remains at high levels throughout the cell cycle in HL-60 cells. DNA topoisomerase III β exists as three isoforms, namely β 1, β 2, and β 3, also produced by alternative splicing. DNA topoisomerase III β 1 is expressed in testes, heart, and skeletal muscle, whereas β 2 is expressed in thymus, kidney, and pancreas.

REFERENCES

1. D-Arpa, P., et al. 1988. cDNA cloning of human DNA topoisomerase I: catalytic activity of a 67.7 kDa carboxyl-terminal fragment. *Proc. Natl. Acad. Sci. USA* 85: 2543-2547.
2. Kunze, N., et al. 1991. Structure of the human type I DNA topoisomerase gene. *J. Biol. Chem.* 266: 9610-9616.
3. Hanai, R., et al. 1996. Human TOP3: a single-copy gene encoding DNA topoisomerase III. *Proc. Natl. Acad. Sci. USA* 93: 3653-3657.
4. Kawasaki, K., et al. 1997. One-megabase sequence analysis of the human immunoglobulin λ gene locus. *Genome Res.* 7: 250-261.
5. Ng, S.W., et al. 1999. A new human topoisomerase III that interacts with SGS1 protein. *Nucleic Acids Res.* 27: 993-1000.

CHROMOSOMAL LOCATION

Genetic locus: TOP3B (human) mapping to 22q11.22; Top3b (mouse) mapping to 16 A3.

SOURCE

Topo III β -1,2,3 (D-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Topo III β -1,2,3 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-11264 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4°C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

Topo III β -1,2,3 (D-16) is recommended for detection of Topo III β -1, Topo III β -2 and Topo III β -3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Topo III β -1,2,3 (D-16) is also recommended for detection of Topo III β -1, Topo III β -2 and Topo III β -3 in additional species, including equine, canine, bovine, porcine and avian.

Positive Controls: Topo III β (m): 293T Lysate: sc-124216.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Topo III β -1,2,3 (D-16): sc-11264. Western blot analysis of Topo III β expression in non-transfected: sc-117752 (A) and mouse Topo III β transfected: sc-124216 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.