

ZAG (H-123): sc-11358

BACKGROUND

ZAG (Zn- α 2-glycoprotein, also designated Zn- α 2-gp) is a soluble, secreted protein found in serum and other body fluids (such as cerebrospinal fluid, blood plasma, urine and sweat). ZAG has a tendency to precipitate with zinc salts, has electrophoretic mobility in the region of the two globulins, and has 18% carbohydrate content. A member of the immunoglobulin superfamily, ZAG has a high degree of sequence similarity to class-I major histocompatibility complex (MHC) antigens. The ZAG structure includes a large groove analogous to class I MHC peptide binding grooves. The crystal structure of ZAG resembles a class I MHC heavy chain but does not bind the class I light chain β -2-Microglobulin, unlike other MHC related proteins. ZAG stimulates lipid degradation in adipocytes and its overexpression causes the extensive fat losses associated with some advanced cancers.

REFERENCES

- Jirka, M., et al. 1973. Zn- α 2-glycoprotein in sweat. *Cas. Lek. Cesk.* 112: 1606-1608.
- Ekman, R., et al. 1976. Renal handling of Zn- α 2-glycoprotein as compared with that of albumin and the retinol-binding protein. *J. Clin. Invest.* 57: 945-954.
- Shibata, S., et al. 1982. Nephritogenic glycoprotein. IX. Plasma Zn- α 2-glycoprotein as a second source of nephritogenic glycoprotein in urine. *Nephron* 31: 170-176.
- Uria, J.A., et al. 1996. Alternative splicing gives rise to two novel long isoforms of Zn- α 2-glycoprotein, a member of the immunoglobulin superfamily. *Gene* 169: 233-236.

CHROMOSOMAL LOCATION

Genetic locus: AZGP1 (human) mapping to 7q22.1; Azgp1 (mouse) mapping to 5 G2.

SOURCE

ZAG (H-123) is a rabbit polyclonal antibody raised against amino acids 13-135 mapping near the N-terminus of ZAG of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

ZAG (H-123) is recommended for detection of ZAG of human and, to a lesser extent, mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for ZAG siRNA (h): sc-36865, ZAG siRNA (m): sc-36866, ZAG shRNA Plasmid (h): sc-36865-SH, ZAG shRNA Plasmid (m): sc-36866-SH, ZAG shRNA (h) Lentiviral Particles: sc-36865-V and ZAG shRNA (m) Lentiviral Particles: sc-36866-V.


Molecular Weight of ZAG: 47 kDa.

Positive Controls: ZAG (h): 293T Lysate: sc-114991.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



ZAG (H-123): sc-11358. Western blot analysis of ZAG expression in non-transfected: sc-117752 (A) and human ZAG transfected: sc-114991 (B) 293T whole cell lysates.

SELECT PRODUCT CITATIONS

- Ding, Z., et al. 2007. Identification of sperm forward motility-related proteins in human seminal plasma. *Mol. Reprod. Dev.* 74: 1124-1131.

MONOS
Satisfaction
Guaranteed

Try **ZAG (1D4): sc-13585** or **ZAG (F-9): sc-365850**, our highly recommended monoclonal alternatives to ZAG (H-123).