

# Nibrin (H-300): sc-11431

## BACKGROUND

DNA repair proteins are necessary for the maintenance of chromosome integrity and are involved in the elimination of premutagenic lesions from DNA. The DNA repair proteins Rad51 and Rad52 are key components of the double-strand-break repair (DSBR) pathway. Rad51 is essential for mitotic and meiotic recombination, and its mutation in yeast and mammalian cells results in chromosome loss. Overexpression of Rad52 confers resistance to ionizing radiation and induces homologous intrachromosomal recombination. Rad52 is thought to be involved in an early stage of Rad51-mediated recombination. Additional proteins involved in the pathway include Dmc1 and nibrin. Dmc1 is specifically involved in meiotic recombination. Nibrin, which complexes with Mre11 and Rad50, is absent in Nijmegen breakage syndrome (NBS) patients.

## CHROMOSOMAL LOCATION

Genetic locus: NBN (human) mapping to 8q21.3; Nbn (mouse) mapping to 4 A2.

## SOURCE

Nibrin (H-300) is a rabbit polyclonal antibody raised against amino acids 1-300 mapping at the N-terminus of Nibrin (NBS1) of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## APPLICATIONS

Nibrin (H-300) is recommended for detection of Nibrin of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Nibrin (H-300) is also recommended for detection of Nibrin in additional species, including equine and bovine.

Suitable for use as control antibody for Nibrin siRNA (h): sc-36061, Nibrin siRNA (m): sc-36062, Nibrin shRNA Plasmid (h): sc-36061-SH, Nibrin shRNA Plasmid (m): sc-36062-SH, Nibrin shRNA (h) Lentiviral Particles: sc-36061-V and Nibrin shRNA (m) Lentiviral Particles: sc-36062-V.

Molecular Weight of Nibrin: 95 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200 or HeLa nuclear extract: sc-2120.


## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2051 or ABC: sc-2018 rabbit IgG Staining Systems.

## DATA



Nibrin (H-300): sc-11431. Immunofluorescence staining of formalin-fixed HeLa cells showing nuclear and cytoplasmic localization. Kindly provided by Yang Xiang, Ph.D., Division of Newborn Medicine, Boston Children's Hospital, Cell Biology Department, Harvard Medical School (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human testis tissue showing nuclear and cytoplasmic staining of cells in seminiferous ducts and Leydig cells (B).

## SELECT PRODUCT CITATIONS

1. Cruet-Hennequart, S., et al. 2008. Enhanced DNA-PK-mediated RPA2 hyperphosphorylation in DNA polymerase  $\eta$ -deficient human cells treated with cisplatin and oxaliplatin. *DNA Repair* 7: 582-596.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.


 MONOS  
 Satisfaction  
 Guaranteed

Try **Nibrin (B-5): sc-515069** or **Nibrin (A-2): sc-374168**, our highly recommended monoclonal alternatives to Nibrin (H-300).