

MAGI-1 (T-17): sc-11524

BACKGROUND

The membrane-associated guanylate kinase (MAGUK) proteins are concentrated at the membrane-cytoskeletal interface where they facilitate the assembly of multiprotein complexes on the inner surface of the plasma membrane. Three protein-protein interaction modules characteristically define MAGUK related proteins: the PDZ domain, the SH3 domain and the guanylate kinase (GuK) domain. The closely related MAGUK proteins, MAGI-1, MAGI-2 and MAGI-3 (membrane associated guanylate kinase inverted-1 and 2), likewise contain the GuK domain and five PDZ domains; however, the SH3 domain is replaced with a WW domain. The transcripts of MAGI-1 are alternatively spliced to produce three distinct proteins having unique carboxy-terminals. Two variants, MAGI-1 α and MAGI-1 β , are associated with the membrane and cytosolic fractions and are primarily expressed in the brain. The third isoform, MAGI-1c, encodes for a nuclear localization signal that localizes MAGI-1c to the nucleus, and it is primarily expressed in the liver and kidney. MAGI-2 and MAGI-3 are localized to the plasma membrane, and they contribute to protein scaffolding by associating with the protein phosphatase PTEN.

REFERENCES

- Anderson, J.M. 1996. Cell signalling: MAGUK magic. *Curr. Biol.* 6: 382-384.
- Dobrosotskaya, I., Guy, R.K. and James, G.L. 1997. MAGI-1, a membrane-associated guanylate kinase with a unique arrangement of protein-protein interaction domains. *J. Biol. Chem.* 272: 31589-31597.
- Wood, J.D., Yuan, J., Margolis, R.L., Colomer, V., Duan, K., Kushi, J., Kaminsky, Z., Kleiderlein, J.J., Sharp, A.H. and Ross, C.A. 1998. Atrophin-1, the DRPLA gene product, interacts with two families of WW domain-containing proteins. *Mol. Cell. Neurosci.* 11: 149-160.
- Dimitratos, S.D., Woods, D.F., Stathakis, D.G. and Bryant, P.J. 1999. Signaling pathways are focused at specialized regions of the plasma membrane by scaffolding proteins of the MAGUK family. *Bioessays* 21: 912-921.
- Dobrosotskaya, I.Y. and James, G.L. 2000. MAGI-1 interacts with β -catenin and is associated with cell-cell adhesion structures. *Biochem. Biophys. Res. Commun.* 270: 903-909.
- Wu, Y., Dowbenko, D., Spencer, S., Laura, R., Lee, J., Gu, Q. and Lasky, L.A. 2000. Interaction of the tumor suppressor PTEN/MMAC with a PDZ domain of MAGI 3, a novel membrane-associated guanylate kinase. *J. Biol. Chem.* 275: 21477-21485
- Wu, X., Hepner, K., Castelino-Prabhu, S., Do, D., Kaye, M.B., Yuan, X.J., Wood, J., Ross, C., Sawyers, C.L. and Whang, Y.E. 2000. Evidence for regulation of the PTEN tumor suppressor by a membrane-localized multi-PDZ domain containing scaffold protein MAGI-2. *Proc. Natl. Acad. Sci. USA* 97: 4233-4238.
- Sakurai, A., Fukuhara, S., Yamagishi, A., Sako, K., Kamioka, Y., Masuda, M., Nakaoka, Y. and Mochizuki, N. 2006. MAGI-1 is required for Rap1 activation upon cell-cell contact and for enhancement of vascular endothelial cadherin-mediated cell adhesion. *Mol. Biol. Cell* 17: 966-976.

CHROMOSOMAL LOCATION

Genetic locus: MAGI1 (human) mapping to 3p14.1.

SOURCE

MAGI-1 (T-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of MAGI-1 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-11524 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

MAGI-1 (T-17) is recommended for detection of MAGI-1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

MAGI-1 (T-17) is also recommended for detection of MAGI-1 in additional species, including canine, bovine and porcine.

Suitable for use as control antibody for MAGI-1 siRNA (h): sc-41999, MAGI-1 shRNA Plasmid (h): sc-41999-SH and MAGI-1 shRNA (h) Lentiviral Particles: sc-41999-V.

Molecular Weight of MAGI-1: 165 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.


 MONOS
 Satisfation
 Guaranteed

Try **MAGI-1 (SS-5): sc-100326**, our highly recommended monoclonal alternative to MAGI-1 (T-17).