

DcR3 (C-15): sc-11643

BACKGROUND

Tumor necrosis factor (TNF) is a pleiotropic cytokine whose function is mediated by two distinct cell surface receptors, designated TNF-R1 and TNF-R2, which are expressed on most cell types. TNF function is primarily mediated through TNF-R1 signaling. Both TNF-R1 and TNF-R2 belong to the growing TNF receptor superfamily which includes FAS antigen and CD40. TNF-R1 contains a cytoplasmic motif, termed the "death domain", that has been found to be necessary for the transduction of the apoptotic signal. The death domain is also found in several other receptors, including FAS, DR2 (or TRUNDD), DR3 (death receptor 3), DR4, DR5, and DR6. TRUNDD, DR4 and DR5 are receptors for the apoptosis-inducing cytokine TRAIL. Non-death domain-containing receptors, designated decoy receptors (DcR1 or TRID, DcR2, and DcR3), associate with specific ligands and may play a role in cellular resistance to apoptotic stimuli.

REFERENCES

1. Tartaglia, L.A., et al. 1993. A novel domain within the 55 kd TNF receptor signals cell death. *Cell* 74: 845-853.
2. Smith, C.A., et al. 1994. The TNF receptor superfamily of cellular and viral proteins: activation, costimulation, and death. *Cell* 76: 959-962.

CHROMOSOMAL LOCATION

Genetic locus: TNFRSF6B (human) mapping to 20q13.33.

SOURCE

DcR3 (C-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of DcR3 (decoy receptor 3) of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-11643 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

DcR3 (C-15) is recommended for detection of DcR3 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for DcR3 siRNA (h): sc-40236, DcR3 shRNA Plasmid (h): sc-40236-SH and DcR3 shRNA (h) Lentiviral Particles: sc-40236-V.


Molecular Weight of DcR3: 33 kDa.

Positive Controls: COLO 320DM cell lysate: sc-2226.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

DATA



DcR3 (C-15): sc-11643. Western blot analysis of human recombinant DcR3 fusion protein.



DcR3 (C-15): sc-11643. Immunoperoxidase staining of formalin fixed, paraffin-embedded human cerebral cortex tissue showing neuropil staining.

SELECT PRODUCT CITATIONS

1. Liu, J., et al. 2015. The expression of death decoy receptor 3 was increased in the patients with primary Sjögren's syndrome. *Clin. Rheumatol.* 34: 879-885.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **DcR3 (F-4): sc-365755** or **DcR3 (A-9): sc-398892**, our highly recommended monoclonal alternatives to DcR3 (C-15).