

Synaptojanin 2 (M-19): sc-12143

BACKGROUND

The inositol polyphosphate 5-phosphatases selectively remove the phosphate from the 5-position of various phosphatidylinositols, which generate second messengers in response to extracellular signals. Synaptojanins are characterized by an N-terminal SAC1-like sequence, a central 5-phosphate domain, and a unique C-terminal sequence and have been shown to use phosphatidylinositol 4,5-bisphosphate as a substrate. Synaptojanins exist as two isoforms, synaptojanin 1 and 2, which differ in the C-terminal domain, and each isoform has multiple variants produced by alternative splicing. Synaptojanin 1 is expressed as two major forms: the shorter is found in brain while the longer is expressed in peripheral tissues. Eight splice variants of synaptojanin 2 have been detected, including a brain specific isoform. Synaptojanins are thought to participate in the endocytosis of synaptic vesicles and the regulation of the actin cytoskeleton.

REFERENCES

- Mitchell, C.A., et al. 1996. Regulation of second messengers by the inositol polyphosphate 5-phosphatases. *Biochem. Soc. Trans.* 24: 994-1000.
- Nemoto, Y., et al. 1997. Synaptojanin 2, a novel synaptojanin isoform with a distinct targeting domain and expression pattern. *J. Biol. Chem.* 272: 30817-30821.
- Zhang, X., et al. 1998. Phosphatidylinositol signalling reactions. *Semin. Cell Dev. Biol.* 9: 153-160.
- Erneux, C., et al. 1998. The diversity and possible functions of the inositol polyphosphate 5-phosphatases. *Biochim. Biophys. Acta* 1436: 185-199.
- Khovtchev, M. and Sudhof, T.C. 1998. Developmentally regulated alternative splicing in a novel Synaptojanin. *J. Biol. Chem.* 273: 2306-2311.
- Seet, L.F., et al. 1998. Molecular cloning of multiple isoforms of Synaptojanin 2 and assignment of the gene to mouse chromosome 17A2-3.1. *Biochem. Biophys. Res. Commun.* 247: 116-122.

CHROMOSOMAL LOCATION

Genetic locus: Synj2 (mouse) mapping to 17 A1.

SOURCE

Synaptojanin 2 (M-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of Synaptojanin 2 of mouse origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-12143 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Synaptojanin 2 (M-19) is recommended for detection of Synaptojanin 2 isoforms δ and ε of mouse origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Synaptojanin 2 siRNA (m): sc-39081, Synaptojanin 2 shRNA Plasmid (m): sc-39081-SH and Synaptojanin 2 shRNA (m) Lentiviral Particles: sc-39081-V.


Molecular Weight of Synaptojanin 2 splice variants: 160/165 kDa.

Positive Controls: Synaptojanin 2 (m): 293T Lysate: sc-123864 or mouse liver extract: sc-2256.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Synaptojanin 2 (M-19): sc-12143. Western blot analysis of Synaptojanin 2 expression in non-transfected: sc-117752 (A) and mouse Synaptojanin 2 transfected: sc-123864 (B) 293T whole cell lysates and mouse liver tissue extract (C).

RESEARCH USE

For research use only, not for use in diagnostic procedures.



Try **Synaptojanin 2 (E-1): sc-390354** or **Synaptojanin 2 (D-11): sc-390247**, our highly recommended monoclonal alternatives to Synaptojanin 2 (M-19).