

PKR (m): 293T Lysate: sc-122612

BACKGROUND

An interferon-inducible, RNA-dependent protein serine/threonine kinase (PKR) has been described. PKR in earlier literature is variously known as DAI, dsJ, PI kinase, p65, p67 or TIK for the mouse kinase; and p68 or p69 for the human kinase. The PKR kinase substrate is the α subunit of protein synthesis initiation factor eIF-2. Phosphorylation of eIF-2 α on Serine 51 results in inhibition of translation. Molecular cDNA clones have been isolated from both human and mouse cells. The serine/threonine kinase catalytic domains map to the carboxy-terminal half of the protein while the RNA-binding domains are located in the amino-terminal region. Three kinds of regulation of PKR enzymatic activity have been described. These include transcriptional regulation in response to interferon, an autoregulatory mechanism controlling PKR expression at the level of translation and post-translational regulation by RNA mediated autophosphorylation.

REFERENCES

- Hershey, J.W.B. 1989. Protein phosphorylation controls translation rates. *J. Biol. Chem.* 264: 20823-20826.
- Meurs, E., et al. 1990. Molecular cloning and characterization of the human double-stranded RNA-activated protein kinase induced by interferon. *Cell* 62: 379-390.
- Icely, P.L., et al. 1991. TIK, a novel serine/threonine kinase, is recognized by antibodies directed against phosphotyrosine. *J. Biol. Chem.* 266: 16073-16077.
- Thomis, D.C., et al. 1992. Mechanism of interferon action: cDNA structure expression, and regulation of the interferon-induced, RNA-dependent PI/eIF-2 α protein kinase from human cells. *Virology* 188: 33-46.
- McCormack, S.J., et al. 1992. Mechanism of interferon action: identification of a RNA binding domain within the N-terminal region of the human RNA-dependent PI/eIF-2 α protein kinase. *Virology* 188: 47-56.
- Samuel, C.E. 1993. The eIF-2 α protein kinases, regulators of translation in eukaryotes from yeasts to humans. *J. Biol. Chem.* 268: 7603-7606.
- Baier, L.J., et al. 1993. The mouse antiphosphotyrosine immunoreactive kinase, TIK, is indistinguishable from the double-stranded RNA-dependent, interferon-induced protein kinase, PKR. *Nucleic Acids Res.* 21: 4830-4835.
- Tanaka, H., et al. 1994. Mechanism of interferon Action: structure of the mouse PKR gene encoding the interferon-inducible RNA-dependent protein kinase. *Proc. Natl. Acad. Sci. USA* 91: 7995-7999.

CHROMOSOMAL LOCATION

Genetic locus: Eif2ak2 (mouse) mapping to 17 E3.

PRODUCT

PKR (m): 293T Lysate represents a lysate of mouse PKR transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

PKR (m): 293T Lysate is suitable as a Western Blotting positive control for mouse reactive PKR antibodies. Recommended use: 10-20 μ l per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

PKR (B-10): sc-6282 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse PKR expression in PKR transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



PKR (B-10) HRP: sc-6282 HRP. Direct western blot analysis of PKR expression in non-transfected: sc-117752 (A) and mouse PKR transfected: sc-122612 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.