

CRF-RI (F-20): sc-12382

BACKGROUND

Individuals suffering from Alzheimer's disease (AD) exhibit dramatic reductions in the content of corticotropin-releasing factor (CRF), increased expression of CRF receptors (CRFRs) and abnormalities in neuronal morphology in affected brain areas. In addition, AD patients show decreased concentrations of CRF in their cerebrospinal fluid, which may contribute to their cognitive impairment. A high affinity CRF binding protein, designated CRF-BP, has been discovered in postmortem brain samples from AD patients. CRF-BP serves to bind and inactivate CRF, reducing the pool of "free CRF" available to bind CRFRs. Two CRF receptors, designated CRF-RI and CRF-RII, exhibit distinct brain localizations. Two forms of CRF-RII, designated CRF-RII α and CRF-RII β , result from alternative mRNA splicing. Urocortin, an additional member of the CRF family, shares 63% sequence identity with urotensin and 45% sequence identity with CRF. Urocortin specifically binds to and activates CRF-RI and CRF-RII, but binds to CRF-RII more efficiently than CRF, suggesting that it may be the true, high affinity ligand for the CRF receptor type II.

REFERENCES

- Behan, D.P., et al. 1995. Displacement of corticotropin releasing factor from its binding protein as a possible treatment for Alzheimer's disease. *Nature* 378: 284-287.
- Behan, D.P., et al. 1995. Corticotropin releasing factor binding protein (CRF-BP) is expressed in neuronal and astrocytic cells. *Brain Res.* 698: 259-264.
- Behan, D.P., et al. 1995. Corticotropin releasing factor (CRF) binding protein: a novel regulator of CRF and related peptides. *Front. Neuroendocrinol.* 16: 362-382.
- Chalmers, D.T., et al. 1995. Localization of novel corticotropin-releasing factor receptor (CRF2) mRNA expression to specific subcortical nuclei in rat brain: comparison with CRF1 receptor mRNA expression. *J. Neurosci.* 15: 6340-6350.
- Lovenberg, T.W., et al. 1995. CRF2 α and CRF2 β receptor mRNAs are differentially distributed between the rat central nervous system and peripheral tissues. *Endocrinology* 136: 4139-4142.
- Liaw, C.W., et al. 1996. Cloning and characterization of the human corticotropin-releasing factor-2 receptor complementary deoxyribonucleic acid. *Endocrinology* 137: 72-77.

CHROMOSOMAL LOCATION

Genetic locus: CRHR1 (human) mapping to 17q21.31.

SOURCE

CRF-RI (F-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of CRF-RI of human origin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-12382 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

CRF-RI (F-20) is recommended for detection of CRF-RI of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for CRF-RI siRNA (h): sc-39914, CRF-RI shRNA Plasmid (h): sc-39914-SH and CRF-RI shRNA (h) Lentiviral Particles: sc-39914-V.

Molecular Weight of CRF-RI: 53-66 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

SELECT PRODUCT CITATIONS

- Yata, A., et al. 2009. Suppression of progesterone production by stress-copin/urocortin 3 in cultured human granulosa-lutein cells. *Hum. Reprod.* 24: 1748-1753.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.