

Synaptotagmin II (S-15): sc-12465

BACKGROUND

Synaptotagmins are a large gene family of synaptic vesicle type III integral membrane proteins that function as regulators of both exocytosis and endocytosis and are involved in neurotransmitter secretion from small secretory vesicles. Calcium binds to Synaptotagmin I which triggers neurotransmitter release at the synapse. Synaptotagmin II is phosphorylated by WNK1 in a process that regulates calcium-dependent interactions. Synaptotagmin III is involved in calcium-dependent exocytosis of secretory vesicles in endocrine cells and neurons. Synaptotagmin IV is expressed in neuronal tissues, and has the highest mRNA levels in the hippocampus. The proximity of the Synaptotagmin IV gene to markers of several psychiatric disorders suggest an involvement of Synaptotagmin IV in human disease. Synaptotagmin V is a dense-core vesicle-specific protein that regulates a specific type of calcium-regulated secretion. Synaptotagmin VI interacts with adaptor protein-2 in a calcium-independent manner. Synaptotagmin VII is widely expressed in non-neuronal tissues.

CHROMOSOMAL LOCATION

Genetic locus: Syt2 (mouse) mapping to 1 E4.

SOURCE

Synaptotagmin II (S-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of Synaptotagmin II of mouse origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-12465 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

Synaptotagmin II (S-15) is recommended for detection of Synaptotagmin II of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Synaptotagmin II (S-15) is also recommended for detection of Synaptotagmin II in additional species, including equine and bovine.

Suitable for use as control antibody for Synaptotagmin II siRNA (m): sc-41313, Synaptotagmin II shRNA Plasmid (m): sc-41313-SH and Synaptotagmin II shRNA (m) Lentiviral Particles: sc-41313-V.


Molecular Weight of Synaptotagmin II: 67 kDa.

Positive Controls: rat brain extract: sc-2392, mouse brain extract: sc-2253 or mouse hypothalamus extract: sc-364242.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Synaptotagmin II (S-15): sc-12465. Western blot analysis of Synaptotagmin II expression in mouse hypothalamus (A), mouse cerebellum (B) and mouse brain (C) tissue extracts.

SELECT PRODUCT CITATIONS

- Atiya-Nasagi, Y., et al. 2005. O-glycosylation is essential for intracellular targeting of Synaptotagmins I and II in non-neuronal specialized secretory cells. *J. Cell Sci.* 118: 1363-1372.
- Chen, G.Y., et al. 2005. Abnormalities caused by carbohydrate alterations in Iβ6-N-acetylglucosaminyltransferase-deficient mice. *Mol. Cell. Biol.* 25: 7828-7838.
- Han, C., et al. 2009. Human SCAMP5, a novel secretory carrier membrane protein, facilitates calcium-triggered cytokine secretion by interaction with SNARE machinery. *J. Immunol.* 182: 2986-2996.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **Synaptotagmin II (26): sc-136089**, our highly recommended monoclonal alternative to Synaptotagmin II (S-15).