

IL-3 (M-20): sc-1259

BACKGROUND

Interleukin-3, or IL-3, is a pleiotropic cytokine that is primarily secreted by activated T lymphocytes and stimulates the proliferation and differentiation of hematopoietic cells. IL-3 not only supports growth of both pluripotent stem cells and the more differentiated committed progenitors, but it also stimulates the functional activity of some fully differentiated cells. IL-3 has also been shown to protect mast cells from undergoing apoptosis. IL-3 exerts its biological effects through a receptor which consists of a ligand-specific α subunit and a signal transducing β subunit common to the IL-3/IL-5/GM-CSF receptors. The carboxy terminus of the β subunit has been shown to be necessary for activation of the MAP kinase signaling pathway. Although the IL-3 receptor has no intrinsic kinase activity, stimulation with IL-3 leads to tyrosine phosphorylation of the JAK/Tyk 2 family member, JAK2, which in turn activates and causes nuclear translocation of Stat5a and Stat5b.

REFERENCES

1. Ihle, J.N., et al. 1985. Interleukin 3. *Methods Enzymol.* 116: 540-552.
2. Yang, Y.C., et al. 1986. Human IL-3 (multi-CSF): identification by expression cloning of a novel hematopoietic growth factor related to murine IL-3. *Cell* 47: 3-10.
3. Mekori, Y.A., et al. 1993. IL-3-dependent murine mast cells undergo apoptosis on removal of IL-3. Prevention of apoptosis by c-kit ligand. *J. Immunol.* 151: 3775-3784.
4. Magnelli, L., et al. 1993. Apoptosis induction in 32D cells by IL-3 withdrawal is preceded by a drop in the intracellular calcium level. *Biochem. Biophys. Res. Comm.* 194: 1394-1397.
5. Mui, A.L.F., et al. 1995. Interleukin-3, granulocyte-macrophage colony stimulating factor and interleukin-5 transduce signals through two STAT5 homologs. *EMBO J.* 14: 1166-1175.
6. Kinoshita, T., et al. 1995. Suppression of apoptotic death in hematopoietic cells by signalling through the IL-3/GM-CSF receptors. *EMBO J.* 14: 266-275.
7. Bagley, C.J., et al. 1995. Interaction of GM-CSF and IL-3 with the common β -chain of their receptors. *J. Leukoc. Biol.* 57: 739-746.

CHROMOSOMAL LOCATION

Genetic locus: IL3 (human) mapping to 5q31.1; Il3 (mouse) mapping to 11 B1.

SOURCE

IL-3 (M-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of IL-3 of mouse origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-1259 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

IL-3 (M-20) is recommended for detection of IL-3 of mouse and, to a lesser extent, rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Suitable for use as control antibody for IL-3 siRNA (m): sc-39622, IL-3 shRNA Plasmid (m): sc-39622-SH and IL-3 shRNA (m) Lentiviral Particles: sc-39622-V.

Molecular Weight of IL-3: 15 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



IL-3 (M-20): sc-1259. Western blot analysis of mouse recombinant IL-3.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **IL-3 (G-11): sc-398210** or **IL-3 (F-6): sc-398177**, our highly recommended monoclonal alternatives to IL-3 (M-20).