

IL-1 β (B122): sc-12742

BACKGROUND

Two forms of interleukin-1, designated IL-1 α and IL-1 β , have been described. Although encoded by distinct genes and exhibiting roughly only 25% sequence identity, IL-1 α and IL-1 β bind to the same receptor and seem to elicit similar biological responses. IL-1 production is generally thought to be associated with inflammation, but it has also been shown to be expressed during kidney development, thymocyte differentiation and cartilage degradation. IL-1 plays a critical role in the regulation of immune response and inflammation, acting as an activator of T and B lymphocytes and natural killer (NK) cells. In T cells, IL-1 stimulates the production of IL-2 and selectively inhibits IL-4 expression. IL-1 induces B cell proliferation and maturation, and immunoglobulin synthesis. NK cells require IL-1 β for production of the anti-pathogen IFN- γ . IL-1 has also been implicated in several pathological conditions including rheumatoid arthritis, inflammatory bowel disease and atherosclerosis.

CHROMOSOMAL LOCATION

Genetic locus: IL1B (human) mapping to 2q13; IL1b (mouse) mapping to 2 F1.

SOURCE

IL-1 β (B122) is an Armenian hamster monoclonal antibody raised against the mature form IL-1 β of mouse origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available azide-free for neutralization, sc-12742 L, 200 μ g/0.1 ml.

IL-1 β (B122) is available conjugated to agarose (sc-12742 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-12742 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-12742 PE), fluorescein (sc-12742 FITC), Alexa Fluor[®] 488 (sc-12742 AF488), Alexa Fluor[®] 546 (sc-12742 AF546), Alexa Fluor[®] 594 (sc-12742 AF594) or Alexa Fluor[®] 647 (sc-12742 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-12742 AF680) or Alexa Fluor[®] 790 (sc-12742 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

IL-1 β (B122) is recommended for detection of IL-1 β precursor and mature forms of mouse, rat, human and hamster origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for IL-1 β siRNA (h): sc-39615, IL-1 β siRNA (m): sc-39616, IL-1 β siRNA (r): sc-45995, IL-1 β shRNA Plasmid (h): sc-39615-SH, IL-1 β shRNA Plasmid (m): sc-39616-SH, IL-1 β shRNA Plasmid (r): sc-45995-SH, IL-1 β shRNA (h) Lentiviral Particles: sc-39615-V, IL-1 β shRNA (m) Lentiviral Particles: sc-39616-V and IL-1 β shRNA (r) Lentiviral Particles: sc-45995-V.


Molecular Weight of IL-1 β precursor: 31 kDa.

Molecular Weight of mature IL-1 β : 17 kDa.


STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

DATA



IL-1 β (B122): sc-12742. Western blot analysis of mouse recombinant IL-1 β .



IL-1 β (B122) HRP: sc-12742 HRP. Direct western blot analysis of mouse recombinant IL-1 β .

SELECT PRODUCT CITATIONS

- O'Shea, E., et al. 2005. 3,4-Methylenedioxymethamphetamine increases pro-interleukin-1 β production and caspase-1 protease activity in frontal cortex, but not in hypothalamus, of dark agouti rats: role of interleukin-1 β in neurotoxicity. *Neuroscience* 135: 1095-1105.
- Leiser, S.F. and Miller, R.A. 2010. Nrf2 signaling, a mechanism for cellular stress resistance in long-lived mice. *Mol. Cell. Biol.* 30: 871-884.
- Jonker, M.A., et al. 2012. Small intestine mucosal immune system response to injury and the impact of parenteral nutrition. *Surgery* 151: 278-286.
- Xue, Z.W., et al. 2013. Effects of Shenshao decoction on the inflammatory response in the aorta of a rat atherosclerotic model. *Chin. J. Integr. Med.* 19: 347-352.
- Di Paola, R., et al. 2016. Ultramicrosized palmitoylethanolamide (PEA-um[®]) in the treatment of idiopathic pulmonary fibrosis. *Pharmacol. Res.* 111: 405-412.
- Prema, A., et al. 2017. Fenugreek seed powder attenuated aluminum chloride-induced Tau pathology, oxidative stress, and inflammation in a rat model of Alzheimer's disease. *J. Alzheimers Dis.* 60: S209-S220.
- Xie, Y., et al. 2018. Electro-acupuncture stimulation prevents remifentanyl-induced postoperative hyperalgesia by suppressing spinal microglia in rats. *Exp. Ther. Med.* 16: 353-359.
- Zhang, M., et al. 2019. AIM2 promotes non-small-cell lung cancer cell growth through inflammasome-dependent pathway. *J. Cell. Physiol.* 234: 20161-20173.
- Guo, Q., et al. 2020. Decreased Jagged1 expression in vascular smooth muscle cells delays endothelial regeneration in arteriovenous graft. *Cardiovasc. Res.* E-published.

RESEARCH USE

For research use only, not for use in diagnostic procedures.