

p-Rb (Ser 780): sc-12901

BACKGROUND

Pediatric cancer retinoblastoma and the formation of other human tumors can be attributed to mutations in the retinoblastoma tumor suppressor gene (Rb). The Rb protein regulates differentiation, apoptosis and cell cycle control by coordinating the cell cycle at G₁-S with transcriptional machinery. During G₁, cyclin D-dependent kinase-mediated phosphorylation of Rb at Ser 795 marks the conversion of Rb from a transcriptionally repressive, hypophosphorylated state to an inactive, phosphorylated state, which may be sustained through mitosis by differential phosphorylation of up to 16 putative serine or threonine residues, including Ser 249/Thr 252, Thr 373, Thr 356, Ser 780, Ser 807/Ser 811, and Thr 821/Thr 826. Hypophosphorylated Rb represses the transcription of genes controlling the cell cycle through direct protein-protein interactions and through the recruitment of histone deacetylase.

CHROMOSOMAL LOCATION

Genetic locus: RB1 (human) mapping to 13q14.2; Rb1 (mouse) mapping to 14 D3.

SOURCE

p-Rb (Ser 780) is available as either goat (sc-12901) or rabbit (sc-12901-R) polyclonal affinity purified antibody raised against a short amino acid sequence containing Ser 780 phosphorylated Rb of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with <0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-12901 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

p-Rb (Ser 780)-R is recommended for detection of Ser 780 phosphorylated Rb of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

p-Rb (Ser 780)-R is also recommended for detection of correspondingly phosphorylated Rb in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for Rb siRNA (h): sc-29468, Rb siRNA (m): sc-29469, Rb shRNA Plasmid (h): sc-29468-SH, Rb shRNA Plasmid (m): sc-29469-SH, Rb shRNA (h) Lentiviral Particles: sc-29468-V and Rb shRNA (m) Lentiviral Particles: sc-29469-V.

Molecular Weight (predicted) of p-Rb: 106 kDa.

Molecular Weight (observed) of p-Rb: 107-140 kDa.


STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.


RESEARCH USE

For research use only, not for use in diagnostic procedures.

DATA



Western blot analysis of Rb phosphorylation in non-transfected: sc-117752 (A,D), untreated human Rb transfected: sc-114014 (B,E) and lambda protein phosphatase (sc-200312A) treated human Rb transfected: sc-114014 (C,F) 293T whole cell lysates. Antibodies tested include p-Rb (Ser 780)-R: sc-12901-R (A,B,C) and Rb (M-153): sc-7905 (D,E,F).



Western blot analysis of non-phosphorylated (A,C) and ERK 2-phosphorylated (B,D) mouse recombinant Rb fusion protein. Antibodies tested include Rb (M-153): sc-7905 (A,B) and p-Rb (Ser 780)-R: sc-12901-R (C,D).

SELECT PRODUCT CITATIONS

- Schmidt, M., et al. 2002. Cell cycle inhibition by FoxO forkhead transcription factors involves downregulation of cyclin D. *Mol. Cell. Biol.* 22: 7842-7852.
- Herrero-González, S., et al. 2009. Connexin43 is involved in the effect of endothelin-1 on astrocyte proliferation and glucose uptake. *Glia* 57: 222-233.
- Li, X., et al. 2009. p27^{KIP1} regulates neurogenesis in the rostral migratory stream and olfactory bulb of the postnatal mouse. *J. Neurosci.* 29: 2902-2914.
- Liu, P., et al. 2009. Replication licensing promotes cyclin D1 expression and G₁ progression in untransformed human cells. *Cell Cycle* 8: 125-136.
- Chen, W., et al. 2010. Cryptotanshinone inhibits cancer cell proliferation by suppressing mammalian target of rapamycin-mediated cyclin D1 expression and Rb phosphorylation. *Cancer Prev. Res.* 3: 1015-1025.
- Sherrill, J.D., et al. 2010. Developmental exposures of male rats to soy isoflavones impact Leydig cell differentiation. *Biol. Reprod.* 83: 488-501.
- Lee, T.J., et al. 2010. Stochastic E2F activation and reconciliation of phenomenological cell-cycle models. *PLoS Biol.* 8 pii: e1000488.
- Zhang, L., et al. 2010. microRNA-141 is involved in a nasopharyngeal carcinoma-related genes network. *Carcinogenesis* 31: 559-566.
- Bianco, M.R., et al. 2011. Cross-talk between cell cycle induction and mitochondrial dysfunction during oxidative stress and nerve growth factor withdrawal in differentiated PC12 cells. *J. Neurosci. Res.* 89: 1302-1315.
- Sarkar, T.R., et al. 2012. Identification of a Src tyrosine kinase/SIAH2 E3 ubiquitin ligase pathway that regulates C/EBPδ expression and contributes to transformation of breast tumor cells. *Mol. Cell. Biol.* 32: 320-332.
- Kim, Y.C., et al. 2012. Angiotensin II regulates activation of Bim via Rb/E2F1 during apoptosis: involvement of interaction between AMPKβ1/2 and Cdk4. *Am. J. Physiol. Lung Cell. Mol. Physiol.* 303: L228-L238.