

Sos 1 (h): 293T Lysate: sc-129810

BACKGROUND

The superfamily of GTP-binding proteins, of which Ras proteins are prototypes, has been implicated in a broad range of biological activities. Studies have identified a family of guanine nucleotide-releasing factors (GRFs) that activate Ras in mammalian cells and an "adapter" protein (Sem 5/GRB2) that appears to mediate the interaction of GRFs with activated receptor molecules. Ras-GRF p140 promotes nucleotide exchange on Ras p21s but not on other members of the Ras gene superfamily. In addition, three mammalian homologs of the *Drosophila* Ras-GRF, Son of Sevenless (Sos), have been described. These include two from mouse, m Sos 1 and m Sos 2, and one from human, h Sos. Vav p95 has been reported to function as a GRF in activation of Ras by the T cell receptor and has been reported to have a domain similar to that of Dbl p115, which is a GRF specific for CDC42Hs. Subsequent to activation, Ras appears to interact with Raf, thereby activating the MAP kinase phosphorylation pathway.

REFERENCES

- Lowenstein, E.J., Daly, R.J., Batzer, A.G., Li, W., Margolis, B., Lammers, R., Ullrich, A., Skolnik, E.Y., Bar-Sagi, D. and Schlessinger, J. 1992. The SH2 and SH3 domain-containing protein GRB2 links receptor tyrosine kinases to Ras signaling. *Cell* 40: 431-442.
- Chardin, P., Camonis, J.H., Gale, N.W., Van Aelst, L., Schlessinger, J., Wigler, M.H. and Bar-Sagi, D. 1993. Human Sos 1: a guanine nucleotide exchange factor for Ras that binds to GRB2. *Science* 260: 1338-1343.
- Skolnik, E.Y., Batzer, A., Li, N., Lee, C.H., Lowenstein, E., Mohammadi, M., Margolis, B. and Schlessinger, J. 1993. The function of GRB2 in linking the Insulin receptor to Ras signaling pathways. *Science* 260: 1953-1955.
- Simon, M.A., Dodsen, G.S. and Rubin, G.M. 1993. An SH3-SH2-SH3 protein is required for p21 Ras 1 activation and binds to sevenless and Sos proteins *in vitro*. *Cell* 73: 169-177.
- Egan, S.E., Giddings, B.W., Brooks, M.W., Buday, L., Sizeland, A.M. and Weinburg, R.A. 1993. Association of Sos Ras exchange protein with GRB2 is implicated in tyrosine kinase signal transduction and transformation. *Nature* 363: 45-51.
- Buday, L. and Downward, J. 1993. Epidermal growth factor regulates p21 Ras through the formation of a complex of receptor, GRB2 adaptor protein, and Sos nucleotide exchange factor. *Cell* 73: 611-620.
- Zhang, X., Settleman, J., Kiriakis, J.M., Takeuchi-Suzuki, E., Elledge, S.J., Marshall, M.S., Bruder, J.T., Rapp, U.R. and Avruch, J. 1993. Normal and oncogenic p21 Ras proteins bind to the amino-terminal regulatory domain of c-RAF-1. *Nature* 364: 308-313.

CHROMOSOMAL LOCATION

Genetic locus: SOS1 (human) mapping to 2p22.1.

PRODUCT

Sos 1 (h): 293T Lysate represents a lysate of human Sos 1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

Sos 1 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive Sos 1 antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

Sos 1 (E-11): sc-55528 is recommended as a positive control antibody for Western Blot analysis of enhanced human Sos 1 expression in Sos 1 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



Sos 1 (E-11): sc-55528. Western blot analysis of Sos 1 expression in non-transfected: sc-117752 (A) and human Sos 1 transfected: sc-129810 (B) 293T whole cell lysates.



Sos 1 (A-9): sc-17793. Western blot analysis of Sos 1 expression in non-transfected: sc-117752 (A) and human Sos 1 transfected: sc-129810 (B) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.