



## Fibulin-2 (3B9A11): sc-130082

### BACKGROUND

Fibulin-1 and Fibulin-2 associate with Fibronectin and other extracellular matrix proteins. In bone marrow, Fibulin-1 and Fibulin-2 bind to Fibronectin in the adherent layer. Fibulin-1 expression is stimulated by estrogen in ovarian cancer cell lines and has been suggested as both an agent of metastasis in ovarian cancer cells and as an indicator for predicting cancer risk or aggressiveness in ovarian carcinomas. The mobility of cancer cells may be inhibited with increasing exposure to Fibulin-1. Fibulin-2 binds to the lectin domains of extracellular matrix proteins aggrecan, versican and brevican. Fibulin-2 is abundantly expressed in heart, placenta and ovarian tissue, where it localizes to basement membranes and connective tissue compartments. In mice, differential Fibulin-2 gene expression correlates with the early phase of diabetic kidneys and glomerulosclerosis.

### REFERENCES

1. Argraves, W.S., et al. 1989. Fibulin, a novel protein that interacts with the Fibronectin receptor  $\beta$ -subunit cytoplasmic domain. *Cell* 58: 623-629.
2. Pan, T.C., et al. 1993. Structure and expression of Fibulin-2, a novel extracellular matrix protein with multiple EGF-like repeats and consensus motifs for calcium binding. *J. Cell Biol.* 123: 1269-1277.
3. Zhang, R.Z., et al. 1994. Fibulin-2 (Fbln-2): human cDNA sequence, mRNA expression, and mapping of the gene on human and mouse chromosomes. *Genomics* 22: 425-430.
4. Clinton, G.M., et al. 1996. Estrogens increase the expression of Fibulin-1, an extracellular matrix protein secreted by human ovarian cancer cells. *Proc. Natl. Acad. Sci. USA* 93: 316-320.
5. Roger, P., et al. 1998. Increased immunostaining of Fibulin-1, an estrogen-regulated protein in the stroma of human ovarian epithelial tumors. *Am. J. Pathol.* 153: 1579-1588.
6. Hayashido, Y., et al. 1998. Estradiol and Fibulin-1 inhibit motility of human ovarian- and breast-cancer cells induced by Fibronectin. *Int. J. Cancer* 75: 654-658.
7. Rochefort, H., et al. 1998. Estrogen receptor mediated inhibition of cancer cell invasion and motility: an overview. *Steroid Biochem. Mol. Biol.* 65: 163-168.

### CHROMOSOMAL LOCATION

Genetic locus: FBLN2 (human) mapping to 3p25.1.

### SOURCE

Fibulin-2 (3B9A11) is a mouse monoclonal antibody raised against a recombinant protein corresponding to amino acids 180-440 of Fibulin-2 of human origin.

### PRODUCT

Each vial contains 200  $\mu$ g IgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

### APPLICATIONS

Fibulin-2 (3B9A11) is recommended for detection of Fibulin-2 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Fibulin-2 siRNA (h): sc-43119, Fibulin-2 shRNA Plasmid (h): sc-43119-SH and Fibulin-2 shRNA (h) Lentiviral Particles: sc-43119-V.

Molecular Weight of Fibulin-2: 195 kDa.

Positive Controls: JAR cell lysate: sc-2276, CCD-1064Sk cell lysate: sc-2263 or JEG-3 whole cell lysate: sc-364255.

### RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:  
 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048.

### STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.