

PLEKHG5 (N-15): sc-130100

BACKGROUND

PLEKHG5 (pleckstrin homology domain-containing family G member 5), also known as GEF720 (guanine nucleotide exchange factor 720) or DSMA4, is predominantly expressed in the peripheral nervous system and brain. PLEKHG5 localizes to the cytoplasm, however when cells are stimulated, PLEKHG5 is found near the perinuclear regions. PLEKHG5 contains the highly-conserved DH-PH module which is considered to be the signature motif of the Dbl family of guanine nucleotide exchange factors (GEFs). PLEKHG5 activates the NF κ B signaling pathway and may be involved in the control of neuronal cell differentiation. It has been suggested that mutations in the PLEKHG5 gene may lead to autosomal recessive distal spinal muscular atrophy (DSMA). Four isoforms of PLEKHG5 exists due to alternative splicing events.

REFERENCES

- Hoffman, G.R. and Cerione, R.A. 2002. Signaling to the Rho GTPases: networking with the DH domain. *FEBS Lett.* 513: 85-91.
- Fuentes, E.J., Karnoub, A.E., Booden, M.A., Der, C.J. and Campbell, S.L. 2003. Critical role of the Pleckstrin homology domain in Dbs signaling and growth regulation. *J. Biol. Chem.* 278: 21188-21196.
- Joseph, R.E. and Norris, F.A. 2005. Substrate specificity and recognition is conferred by the Pleckstrin homology domain of the Dbl family guanine nucleotide exchange factor P-Rex2. *J. Biol. Chem.* 280: 27508-27512.
- Baumeister, M.A., Rossman, K.L., Sondek, J. and Lemmon, M.A. 2006. The Dbs PH domain contributes independently to membrane targeting and regulation of guanine nucleotide-exchange activity. *Biochem. J.* 400: 563-572.
- Online Mendelian Inheritance in Man, OMIM[™]. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 611101. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Maystadt, I., Rezsöhazy, R., Barkats, M., Duque, S., Vannuffel, P., Remacle, S., Lambert, B., Najimi, M., Sokal, E., Munnich, A., Viollet, L. and Verellen-Dumoulin, C. 2007. The nuclear factor κ B-activator gene PLEKHG5 is mutated in a form of autosomal recessive lower motor neuron disease with childhood onset. *Am. J. Hum. Genet.* 81: 67-76.
- Rapley, J., Tybulewicz, V.L. and Rittinger, K. 2008. Crucial structural role for the PH and C1 domains of the Vav1 exchange factor. *EMBO Rep.* 9: 655-661.

CHROMOSOMAL LOCATION

Genetic locus: PLEKHG5 (human) mapping to 1p36.31.

SOURCE

PLEKHG5 (N-15) is a purified rabbit polyclonal antibody raised against a peptide mapping at the N-terminus of PLEKHG5 of human origin.

PRODUCT

Each vial contains 100 μ g IgG in 1.0 ml PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

PLEKHG5 (N-15) is recommended for detection of PLEKHG5 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).


Suitable for use as control antibody for PLEKHG5 siRNA (h): sc-88767, PLEKHG5 shRNA Plasmid (h): sc-88767-SH and PLEKHG5 shRNA (h) Lentiviral Particles: sc-88767-V.

Molecular Weight (predicted) of PLEKHG5: 117 kDa.

Molecular Weight (observed) of PLEKHG5: 102 kDa.

Positive Controls: Jurkat whole cell lysate: sc-2204, MCF7 whole cell lysate: sc-2206 or NTERA-2 cl.D1 whole cell lysate: sc-364181.

DATA



PLEKHG5 (N-15): sc-130100. Western blot analysis of PLEKHG5 expression in Jurkat (A), MCF7 (B) and NTERA-2 cl.D1 (C) whole cell lysates.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **PLEKHG5 (5A9): sc-517108**, our highly recommended monoclonal alternative to PLEKHG5 (N-15).