



S-100 α/β chain (S100 217): sc-130110

BACKGROUND

The family of EF-hand type Ca^{2+} -binding proteins includes calbindin (previously designated vitamin D-dependent Ca^{2+} -binding protein), S-100 α and β , calgranulins A (also designated MRP8), B (also designated MRP14) and C (S-100 like proteins), and the parvalbumin family members, including parvalbumin α and parvalbumin β (also designated oncomodulin). The S-100 protein is involved in the regulation of cellular processes such as cell cycle progression and differentiation. Research also indicates that the S-100 protein may function in the activation of Ca^{2+} -induced Ca^{2+} release, inhibition of microtubule assembly and inhibition of protein kinase C mediated phosphorylation. Two S-100 subunits, sharing 60% sequence identity, have been described as S-100 α chain and S-100 β chain. Three S-100 dimeric forms have been characterized, differing in their subunit composition of either two α chains, two β chains or one α and one β chain. S-100 localizes to the cytoplasm and nuclei of astrocytes, Schwann's cells, ependymomas and astroglomas. S-100 is also detected in almost all benign naevi, malignant melanocytic tumours and in Langerhans cells in the skin. Calbindin, S-100 proteins and parvalbumin proteins are each expressed in neural tissues. In addition, S-100 α and β are present in a variety of other tissues, and calbindin is present in intestine and kidney.

REFERENCES

- Pfyster, G.E., Faivre-Bauman, A., Tixier-Vidal, A., Norman, A.W. and Heizmann, C.W. 1987. Developmental and functional studies of parvalbumin and Calbindin D28K in hypothalamic neurons grown in serum-free medium. *J. Neurochem.* 49: 442-451.
- Heizmann, C.W. 1988. Calcium-binding proteins of the EF-type. *J. Cardiovasc. Pharmacol.* 5: S30-S37.
- Kagi, U., Chafouleas, J.G., Norman, A.W. and Heizmann, C.W. 1988. Developmental appearance of the Ca^{2+} -binding proteins parvalbumin, Calbindin D28K, S-100 proteins and calmodulin during testicular development in the rat. *Cell Tissue Res.* 252: 359-365.
- Zimmer, D.B., Song, W. and Zimmer, W.E. 1991. Isolation of a rat S-100 α cDNA and distribution of its mRNA in rat tissues. *Brain Res. Bull.* 27: 157-162.
- Rickmann, M. and Wolff, J.R. 1995. S-100 protein expression in subpopulations of neurons of rat brain. *Neuroscience* 67: 977-991.
- Wang, Y.Z. and Christakos, S. 1995. Retinoic acid regulates the expression of the calcium binding protein, Calbindin D28K. *Mol. Endocrinol.* 9: 1510-1521.
- Muntener, M., Kaser, L., Weber, J. and Berchtold, M.W. 1995. Increase of skeletal muscle relaxation speed by direct injection of parvalbumin cDNA. *Proc. Natl. Acad. Sci. USA* 92: 6504-6508.
- Hitomi, J., Yamaguchi, K., Kikuchi, Y., Kimura, T., Maruyama, K. and Nagasaki, K. 1996. A novel calcium-binding protein in amniotic fluid. CAAF1: its molecular cloning and tissue distribution. *J. Cell. Sci.* 109: 805-815.
- Beaudeau, J., Dequen, L. and Foglietti, M. 1999. Pathophysiologic aspects of S-100 β protein: a new biological marker of brain pathology. *Ann. Biol. Clin.* 57: 261-272.

CHROMOSOMAL LOCATION

Genetic locus: S100A1 (human) mapping to 1q21.3, S100B (human) mapping to 21q22.3; S100a1 (mouse) mapping to 3 F1-F2, S100b (mouse) mapping to 10 C1.

SOURCE

S-100 α/β chain (S100 217) is a mouse monoclonal antibody raised against S-100 of cow origin.

PRODUCT

Each vial contains 100 μg IgG₁ in 1.0 ml PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

S-100 α/β chain (S100 217) is recommended for detection of S-100 α and β chains of mouse, rat, human, cow and monkey origin by immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).

Molecular Weight of S-100 α chain: 11 kDa.

Molecular Weight of S-100 β dimer: 21 kDa.

Molecular Weight of S-100 β chain: 10 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Immunofluorescence: use goat anti-mouse IgG-FITC: sc-2010 (dilution range: 1:100-1:400) or goat anti-mouse IgG-TR: sc-2781 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 2) Immunohistochemistry: use ImmunoCruz™: sc-2050 or ABC: sc-2017 mouse IgG Staining Systems.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.