

# DGK- $\beta$ (C-24): sc-130131

## BACKGROUND

Diacylglycerol kinases (DGKs) phosphorylate diacylglycerol (DAG) to produce phosphatidic acid. DAG and phosphatidic acid are lipids that act as second messengers in signaling cascades. DGK- $\alpha$  influences cell activation and secretion of lethal exosomes, which in turn control cell death. DGK- $\beta$  is abundant in restricted brain regions such as the caudate putamen and olfactory tubercle. DGK- $\gamma$  encodes full-length and truncated transcripts that are present in a range of human tissues, with greatest expression observed in retina. DGK- $\delta$  is most abundant in skeletal muscle. DGK- $\epsilon$  shows specificity for arachidonyl-containing diacylglycerol and is expressed predominantly in testis. DGK- $\zeta$  is most abundant in brain and muscle. DGK- $\eta$  is closely related to DGK- $\delta$ . DGK- $\theta$  is most abundant in the cerebellum and hippocampus. DGK- $\iota$  is present in brain and retina as a predominant transcript of more than 12 kb, including a long 3-prime untranslated region, with additional low abundance transcripts of 9.5 and 7.5 kb. DGKs have structural motifs that play regulatory roles, and these motifs form the basis for dividing the DGKs into five subtypes.

## REFERENCES

1. Schaap, D., et al. 1990. Purification, cDNA-cloning and expression of human diacylglycerol kinase. *FEBS Lett.* 275: 151-158.
2. Goto, K., et al. 1993. Molecular cloning and expression of a 90-kDa diacylglycerol kinase that predominantly localizes in neurons. *Proc. Natl. Acad. Sci. USA* 90: 7598-7602.
3. Masai, I., et al. 1993. *Drosophila* retinal degeneration A gene encodes an eye-specific diacylglycerol kinase with cysteine-rich zinc-finger motifs and ankyrin repeats. *Proc. Natl. Acad. Sci. USA* 90: 11157-11161.
4. Kai, M., et al. 1994. Molecular cloning of a diacylglycerol kinase isozyme predominantly expressed in human retina with a truncated and inactive enzyme expression in most other human cells. *J. Biol. Chem.* 269: 18492-18498.
5. Sakane, F., et al. 1996. Molecular cloning of a novel diacylglycerol kinase isozyme with a pleckstrin homology domain and a C-terminal tail similar to those of the EPH family of protein-tyrosine kinases. *J. Biol. Chem.* 271: 8394-8401.

## CHROMOSOMAL LOCATION

Genetic locus: DGKB (human) mapping to 7p21.2; Dgkb (mouse) mapping to 12 A3.

## SOURCE

DGK- $\beta$  (C-24) is a purified rabbit polyclonal antibody raised against a peptide mapping near the C-terminus of DGK- $\beta$  of human origin.

## PRODUCT

Each vial contains 100  $\mu$ g IgG in 1.0 ml PBS with < 0.1% sodium azide and 0.1% gelatin.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

DGK- $\beta$  (C-24) is recommended for detection of DGK- $\beta$  of mouse and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Suitable for use as control antibody for DGK- $\beta$  siRNA (h): sc-89505, DGK- $\beta$  siRNA (m): sc-155882, DGK- $\beta$  shRNA Plasmid (h): sc-89505-SH, DGK- $\beta$  shRNA Plasmid (m): sc-155882-SH, DGK- $\beta$  shRNA (h) Lentiviral Particles: sc-89505-V and DGK- $\beta$  shRNA (m) Lentiviral Particles: sc-155882-V.

Molecular Weight of DGK- $\beta$ : 91 kDa.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2051 or ABC: sc-2018 rabbit IgG Staining Systems.

## DATA



DGK- $\beta$  (C-24): sc-130131. Western blot analysis of DGK- $\beta$  expression in mouse liver tissue extract.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.


 MONOS  
Satisfaction  
Guaranteed

Try **DGK- $\beta$  (G-11): sc-515090**, our highly recommended monoclonal alternative to DGK- $\beta$  (C-24).