

p-p107 (Ser 975): sc-130209

BACKGROUND

The human retinoblastoma gene product plays an important role in the negative regulation of cell proliferation. Functional inactivation of Rb can be mediated either through mutation or as a consequence of interaction with DNA tumor virus encoded proteins. pRb and the structurally related p107 form complexes with E2F, a transcription factor originally identified through its role in transcriptional activation of the adenovirus E2 promoter. Moreover, pRb and p107 share a high degree of structural homology in the adenovirus E1A binding domain (i.e. "pocket region") that is believed to play a primary role in the function of these proteins. A protein designated p130 shows a high degree of identity with pRb and p107 and also possesses a pocket region. Human p107 is subject to phosphorylation on a variety of residues, including Ser 975, an event which leads to the subsequent inactivation of p107.

REFERENCES

1. Kovcsdi, I., et al. 1986. Identification of a cellular transcription factor involved in E1A transactivation. *Cell* 45: 219-228.
2. Chellappan, S., et al. 1991. The E2F transcription factor is a cellular target for the Rb protein. *Cell* 65: 1053-1061.
3. Chittenden, T., et al. 1991. The T/E1A-binding domain of the retinoblastoma product can interact selectively with a sequence-specific DNA-binding protein. *Cell* 65: 1073-1082.
4. Bandara, L., et al. 1991. Cyclin A and the retinoblastoma gene product complex with a common transcription factor. *Nature* 352: 249-251.
5. Helin, K., et al. 1992. A cDNA encoding a pRb-binding protein with properties of the transcription factor E2F. *Cell* 70: 337-350.
6. Lees, E., et al. 1992. Cyclin E/Cdk2 and cyclin A/Cdk2 kinases associate with p107 and E2F in a temporally distinct manner. *Genes Dev.* 6: 1874-1885.
7. Barbie, T.U., et al. 2003. Mullerian inhibiting substance inhibits cervical cancer cell growth via a pathway involving p130 and p107. *Proc. Natl. Acad. Sci. USA* 100: 15601-15606.

CHROMOSOMAL LOCATION

Genetic locus: RBL1 (human) mapping to 20q11.23.

SOURCE

p-p107 (Ser 975) is a rabbit polyclonal antibody raised against a short amino acid sequence containing Ser 975 phosphorylated p107 of human origin.

PRODUCT

Each vial contains 100 µg IgG in 1.0 ml PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

p-p107 (Ser 975) is recommended for detection of Ser 975 phosphorylated p107 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for p107 siRNA (h): sc-29423, p107 shRNA Plasmid (h): sc-29423-SH and p107 shRNA (h) Lentiviral Particles: sc-29423-V.


Molecular Weight of p-p107 isoforms: 68/121 kDa.

Positive Controls: A2058 whole cell lysate: sc-364178.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto B Blocking Reagent: sc-2335 (use 50 mM NaF, sc-24988, as diluent) and Western Blotting Luminol Reagent: sc-2048 and Lambda Phosphatase: sc-200312A. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2051 or ABC: sc-2018 rabbit IgG Staining Systems.

DATA



p-p107 (Ser 975): sc-130209. Western blot analysis of p-p107 expression in A2058 whole cell lysate.



p-p107 (Ser 975): sc-130209. Immunoperoxidase staining of formalin fixed, paraffin-embedded human cancer tissue showing cytoplasmic staining.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.