

TICAM-1 (C-14): sc-130621

BACKGROUND

Toll/interleukin-1 receptor/resistance (TIR) adaptor protein (Trif, TICAM-1) can physically bind TIR domains and influence cell signaling. TICAM-1 interacts with TLR3 and mediates dsRNA activation of interferon- β through NF κ B, AP1 or IRF-3. Human TICAM-1 maps to chromosome 19p13.3.

REFERENCES

- Yamamoto, M., Sato, S., Mori, K., Hoshino, K., Takeuchi, O., Takeda, K. and Akira, S. 2002. Cutting edge: a novel toll/IL-1 receptor domain-containing adapter that preferentially activates the IFN- β promoter in the toll-like receptor signaling. *J. Immunol.* 169: 6668-6672.
- Hoebe, K., Du, X., Georgel, P., Janssen, E., Tabeta, K., Kim, S.O., Goode, J., Lin, P., Mann, N., Mudd, S., Crozat, K., Sovath, S., Han, J. and Beutler, B. 2003. Identification of Lps2 as a key transducer of MyD88-independent TIR signalling. *Nature* 424: 743-748.
- Oshiumi, H., Matsumoto, M., Funami, K., Akazawa, T. and Seya, T. 2003. TICAM-1, an adaptor molecule that participates in toll-like receptor 3-mediated interferon- β induction. *Nat. Immunol.* 4: 161-167.
- Hardy, M.P., McGettrick, A.F. and O'Neill, L.A. 2004. Transcriptional regulation of the human TRIF (TIR domain-containing adaptor protein inducing interferon- β) gene. *Biochem. J.* 380: 83-93.
- Li, K., Foy, E., Ferrecon, J.C., Nakamura, M., Ferrecon, A.C., Ikeda, M., Ray, S.C., Gale, M. Jr. and Lemon, S.M. 2005. Immune evasion by hepatitis C virus NS3/4A protease-mediated cleavage of the toll-like receptor 3 adaptor protein TRIF. *Proc. Natl. Acad. Sci. USA* 102: 2992-2997.
- Seya, T., Oshiumi, H., Sasai, M., Akazawa, T. and Matsumoto, M. 2005. TICAM-1 and TICAM-2: toll-like receptor adapters that participate in induction of type 1 interferons. *Int. J. Biochem. Cell Biol.* 37: 524-529.
- Sasai, M., Oshiumi, H., Matsumoto, M., Inoue, N., Fujita, F., Nakanishi, M. and Seya T. 2005. Cutting Edge: NF κ B-activating kinase-associated protein 1 participates in TLR3/toll-IL-1 homology domain-containing adapter molecule-1-mediated IFN regulatory factor 3 activation. *J. Immunol.* 174: 27-30.
- LocusLink Report (LocusID: 148022). <http://www.ncbi.nlm.nih.gov/LocusLink>

CHROMOSOMAL LOCATION

Genetic locus: TICAM1 (human) mapping to 19p13.3.

SOURCE

TICAM-1 (C-14) is a purified rabbit polyclonal antibody raised against a 14 amino acid synthetic peptide within the C-terminus region of TICAM-1 of human origin.

PRODUCT

Each vial contains 50 μ g IgG in 500 μ l PBS with < 0.1% sodium azide and 0.1% gelatin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

TICAM-1 (C-14) is recommended for detection of TICAM-1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)].

Suitable for use as control antibody for TICAM-1 siRNA (h): sc-106845, TICAM-1 shRNA Plasmid (h): sc-106845-SH and TICAM-1 shRNA (h) Lentiviral Particles: sc-106845-V.

Molecular Weight of TICAM-1: 66 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **TICAM-1 (E-7): sc-514384**, our highly recommended monoclonal alternative to TICAM-1 (C-14).