

STRAP (3G6): sc-130671

BACKGROUND

Smad proteins play an important role in the intracellular signalling of the TGF β superfamily of extracellular polypeptides. Two Smad proteins, Smad6 and Smad7, function as antagonists to TGF β signalling. STRAP, another antagonist to the TGF β signalling pathway, specifically interacts with Smad7, but not Smad6, to synergistically block TGF β -induced transcriptional activation. The gene encoding the human homolog of STRAP (as designated in mouse), called UNR-interacting protein, maps to chromosome 12p12.3. UNR-interacting protein is 97% homologous to STRAP at the amino acid level. The UNR-interacting protein binds UNR, a cytoplasmic RNA-binding protein with five cold-shock domains that is involved in RNA translation. The presence of the STRAP gene in a variety of species from mammals to yeast, indicates that STRAP function is evolutionarily conserved in eukaryotic cells.

REFERENCES

1. Datta, P.K., Chytil, A., Gorska, A.E. and Moses, H.L. 1998. Identification of STRAP, a novel WD domain protein in transforming growth factor- β signaling. *J. Biol. Chem.* 273: 34671-34674.
2. Hunt, S.L., Hsuan, J.J., Totty, N. and Jackson, R.J. 1999. unr, a cellular cytoplasmic RNA-binding protein with five cold-shock domains, is required for internal initiation of translation of human rhinovirus RNA. *Genes Dev.* 13: 437-448.
3. Datta, P.K. and Moses, H.L. 2000. STRAP and Smad7 synergize in the inhibition of transforming growth factor β signaling. *Mol. Cell. Biol.* 20: 3157-3167.
4. Zhao, J., Shi, W., Chen, H. and Warburton, D. 2000. Smad7 and Smad6 differentially modulate transforming growth factor β induced inhibition of embryonic lung morphogenesis. *J. Biol. Chem.* 275: 23992-23997.
5. Locus Link (LocusID: 11171) <http://www.ncbi.nlm.nih.gov/LocusLink/>

CHROMOSOMAL LOCATION

Genetic locus: STRAP (human) mapping to 12p12.3; Strap (mouse) mapping to 6 G1.

SOURCE

STRAP (3G6) is a mouse monoclonal antibody raised against recombinant STRAP fusion protein of human origin.

PRODUCT

Each vial contains 200 μ g IgG $_1$ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

APPLICATIONS

STRAP (3G6) is recommended for detection of STRAP of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)] and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for STRAP siRNA (h): sc-44129, STRAP siRNA (m): sc-153911, STRAP shRNA Plasmid (h): sc-44129-SH, STRAP shRNA Plasmid (m): sc-153911-SH, STRAP shRNA (h) Lentiviral Particles: sc-44129-V and STRAP shRNA (m) Lentiviral Particles: sc-153911-V.


Molecular Weight of STRAP: 39 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, c4 whole cell lysate: sc-364186 or Neuro-2A whole cell lysate: sc-364185.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



STRAP (3G6): sc-130671. Western blot analysis of STRAP expression in c4 (A), Neuro-2A (B), RPE-J (C) and KNRK (D) whole cell lysates.



STRAP (3G6): sc-130671. Western blot analysis of STRAP expression in HeLa (A), Hep G2 (B) and A549 (C) whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.