

ADCK5 (A-25): sc-130680

BACKGROUND

CABC1 (chaperone activity of bc1 complex-like), also known as COQ8 or ADCK3 (aarF domain-containing protein kinase 3) is a 647 amino acid mitochondrial protein that belongs to the ADCK protein kinase. Ubiquitously expressed with higher expression in heart and skeletal muscle, CABC1 is thought to function as a chaperone in the proper assembly of protein complexes found in the respiratory chain. CABC1 expression is induced both in response to DNA damage and by the tumor suppressor p53. When CABC1 expression is inhibited, p53-induced apoptosis is partially suppressed, suggesting a possible role for CABC1 in tumor suppression. Mutations in the gene encoding CABC1 may be implicated in ubiquinone deficiency which can lead to cerebellar ataxia and seizures. Four isoforms of CABC1 exist due to alternative splicing events. Other members of the ADCK protein kinase family include ADCK1, ADCK2, ADCK4 and ADCK5.

REFERENCES

1. Iizumi, M., Arakawa, H., Mori, T., Ando, A. and Nakamura, Y. 2002. Isolation of a novel gene, CABC1, encoding a mitochondrial protein that is highly homologous to yeast activity of bc1 complex. *Cancer Res.* 62: 1246-1250.
2. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 606980. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
3. Wan, D., Gong, Y., Qin, W., Zhang, P., Li, J., Wei, L., Zhou, X., Li, H., Qiu, X., Zhong, F., He, L., Yu, J., Yao, G., Jiang, H., Qian, L., Yu, Y., Shu, H., Chen, X., Xu, H., Guo, M., Pan, Z., Chen, Y., Ge, C., Yang, S. and Gu, J. 2004. Large-scale cDNA transfection screening for genes related to cancer development and progression. *Proc. Natl. Acad. Sci. USA* 101: 15724-15729.
4. Mollet, J., Delahodde, A., Serre, V., Chretien, D., Schlemmer, D., Lombes, A., Boddaert, N., Desguerre, I., de Lonlay, P., de Baulny, H.O., Munnich, A. and Rötig, A. 2008. CABC1 gene mutations cause ubiquinone deficiency with cerebellar ataxia and seizures. *Am. J. Hum. Genet.* 82: 623-630.
5. Lagier-Tourenne, C., Tazir, M., López, L.C., Quinzii, C.M., Assoum, M., Drouot, N., Busso, C., Makri, S., Ali-Pacha, L., Benhassine, T., Anheim, M., Lynch, D.R., Thibault, C., Plewniak, F., Bianchetti, L., Tranchant, C., Poch, O., et al. 2008. ADCK3, an ancestral kinase, is mutated in a form of recessive ataxia associated with coenzyme Q10 deficiency. *Am. J. Hum. Genet.* 82: 661-672.

CHROMOSOMAL LOCATION

Genetic locus: ADCK5 (human) mapping to 8q24.3; Adck5 (mouse) mapping to 15 D3.

SOURCE

ADCK5 (A-25) is a purified rabbit polyclonal antibody raised against a peptide mapping near the N-terminus of ADCK5 of human origin.

PRODUCT

Each vial contains 100 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

ADCK5 (A-25) is recommended for detection of ADCK5 of mouse and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for ADCK5 siRNA (h): sc-77713, ADCK5 siRNA (m): sc-140877, ADCK5 shRNA Plasmid (h): sc-77713-SH, ADCK5 shRNA Plasmid (m): sc-140877-SH, ADCK5 shRNA (h) Lentiviral Particles: sc-77713-V and ADCK5 shRNA (m) Lentiviral Particles: sc-140877-V.


Molecular Weight of ADCK5: 66 kDa.

Positive Controls: mouse brain extract: sc-2253.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

DATA



ADCK5 (A-25): sc-130680. Western blot analysis of ADCK5 expression in mouse brain tissue extract.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.