

frizzled-1 (V-22): sc-130758

BACKGROUND

The frizzled gene, originally identified in *Drosophila melanogaster*, is involved in the development of tissue polarity. The mammalian homolog of frizzled as well as several secreted mammalian frizzled-related proteins (FRPs) have been described. The frizzled proteins contain seven transmembrane domains, a cysteine-rich domain in the extracellular region and a carboxy terminal Ser/Thr-xxx-Val motif. They function as receptors for Wnt and are generally coupled to G proteins. The frizzled-1 gene is expressed in adult heart, placenta, lung, kidney, pancreas, prostate and ovary and in fetal lung and kidney.

REFERENCES

1. Wang, Y., et al. 1996. A large family of putative transmembrane receptors homologous to the product of the *Drosophila* tissue polarity gene frizzled. *J. Biol. Chem.* 271: 4468-4476.
2. Yang-Snyder, J., et al. 1996. A frizzled homolog functions in a vertebrate Wnt signaling pathway. *Curr. Biol.* 6: 1302-1306.
3. Rattner, A., et al. 1997. A family of secreted proteins contains homology to the cysteine-rich ligand-binding domain of frizzled receptors. *Proc. Natl. Acad. Sci. USA* 94: 2859-2863.
4. Finch, P.W., et al. 1997. Purification and molecular cloning of a secreted, frizzled-related antagonist of Wnt action. *Proc. Natl. Acad. Sci. USA* 94: 6770-6775.

CHROMOSOMAL LOCATION

Genetic locus: FZD1 (human) mapping to 7q21.13.

SOURCE

frizzled-1 (V-22) is a purified rabbit polyclonal antibody raised against a peptide mapping within an internal region of frizzled-1 of human origin.

PRODUCT

Each vial contains 100 µg IgG in 1.0 ml PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

frizzled-1 (V-22) is recommended for detection of frizzled-1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for frizzled-1 siRNA (h): sc-39977, frizzled-1 shRNA Plasmid (h): sc-39977-SH and frizzled-1 shRNA (h) Lentiviral Particles: sc-39977-V.


Molecular Weight of frizzled-1: 71 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2051 or ABC: sc-2018 rabbit IgG Staining Systems.

DATA



frizzled-1 (V-22): sc-130758. Western blot analysis of frizzled-1 expression in HeLa whole cell lysate.



frizzled-1 (V-22): sc-130758. Immunoperoxidase staining of formalin-fixed, paraffin-embedded human prostate carcinoma tissue showing cytoplasmic and membrane localization.

SELECT PRODUCT CITATIONS

1. Liu, H., et al. 2014. Sox9 regulates hyperexpression of Wnt1 and Fzd1 in human osteosarcoma tissues and cells. *Int. J. Clin. Exp. Pathol.* 7: 4795-4805.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



MONOS
Satisfaction
Guaranteed

Try **frizzled-1 (E-7): sc-398082**, our highly recommended monoclonal alternative to frizzled-1 (V-22).