

GCAP3 (W-23): sc-130764

BACKGROUND

The intracellular stimulation of guanylate cyclase (GC) by calcium, a key event in the recovery of the dark state of rod photoreceptors after exposure to light, is mediated by guanylate cyclase-activating proteins (GCAP). GCAPs are calcium-binding proteins belonging to the calmodulin superfamily and are specifically expressed in retina. GCAP3 (guanylyl cyclase-activating protein 3), also known as GUCA1C (guanylate cyclase activator 1C), is a 209 amino acid EF-hand calcium binding protein that is activated by the decrease in calcium from the absorption of light by rhodopsin. Activation of GCAP3 leads to stimulation of guanylate cyclase 1 and 2 (GC1 and GC2), which increases cGMP concentration. Calcium sensitive regulation of GC is essential in recovery of the rod receptor dark state following light exposure. There are two isoforms of GCAP3 that are produced as a result of alternative splicing events.

REFERENCES

- Haeseleer, F., Sokal, I., Li, N., Pettenati, M., Rao, N., Bronson, D., Wechter, R., Baehr, W. and Palczewski, K. 1999. Molecular characterization of a third member of the guanylyl cyclase-activating protein subfamily. *J. Biol. Chem.* 274: 6526-6535.
- Palczewski, K., Verlinde, C.L. and Haeseleer, F. 1999. Molecular mechanism of visual transduction. *Novartis Found. Symp.* 224: 191-204
- Gorczyca, W.A. and Sokal, I. 2002. GCAPs: Ca²⁺-sensitive regulators of retGC. *Adv. Exp. Med. Biol.* 514: 319-332.
- Imanishi, Y., Li, N., Sokal, I., Sowa, M.E., Lichtarge, O., Wensel, T.G., Saperstein, D.A., Baehr, W. and Palczewski, K. 2002. Characterization of retinal guanylate cyclase-activating protein 3 (GCAP3) from zebrafish to man. *Eur. J. Neurosci.* 15: 63-78.
- Palczewski, K., Sokal, I. and Baehr, W. 2004. Guanylate cyclase-activating proteins: structure, function, and diversity. *Biochem. Biophys. Res. Commun.* 322: 1123-1130.
- Imanishi, Y., Yang, L., Sokal, I., Filipek, S., Palczewski, K. and Baehr, W. 2004. Diversity of guanylate cyclase-activating proteins (GCAPs) in teleost fish: characterization of three novel GCAPs (GCAP4, GCAP5, GCAP7) from zebrafish (*Danio rerio*) and prediction of eight GCAPs (GCAP1-8) in pufferfish (*Fugu rubripes*). *J. Mol. Evol.* 59: 204-217.
- Stephen, R., Palczewski, K. and Sousa, M.C. 2006. The crystal structure of GCAP3 suggests molecular mechanism of GCAP-linked cone dystrophies. *J. Mol. Biol.* 359: 266-275.
- Online Mendelian Inheritance in Man, OMIM[™]. 2006. Johns Hopkins University, Baltimore, MD. MIM Number: 605128. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: GUCA1C (human) mapping to 3q13.13.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

SOURCE

GCAP3 (W-23) is a purified rabbit polyclonal antibody raised against a peptide mapping near the N-terminus of GCAP3 of human origin.

PRODUCT

Each vial contains 100 µg IgG in 1.0 ml PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

GCAP3 (W-23) is recommended for detection of GCAP3 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for GCAP3 siRNA (h): sc-75114, GCAP3 shRNA Plasmid (h): sc-75114-SH and GCAP3 shRNA (h) Lentiviral Particles: sc-75114-V.


Molecular Weight of GCAP3: 24 kDa.

Positive Controls: Y79 cell lysate: sc-2240.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

DATA



GCAP3 (W-23): sc-130764. Western blot analysis of GCAP3 expression in Y79 whole cell lysate.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.