

Rsk-1 (W-23): sc-130870

BACKGROUND

The family of ribosomal S6 kinases (Rsk), designated Rsk-1 (or MAPKAP kinase-1), Rsk-2 and Rsk-3, are intracellular serine/threonine kinases that are important signaling intermediates in response to a broad range of ligand-activated receptor tyrosine kinases. A unique feature common to the members of the Rsk family is that each possesses two non-identical complete kinase catalytic domains. An additional Rsk protein, Rsk-4, shows a high level of homology to the three previously isolated members of the human Rsk family. Rsk-4 is most abundantly expressed in brain and kidney and plays a role in normal neuronal development. The family of ribosomal S6 kinases includes p70 S6 kinase and p70 S6 kinase β , which are thought to have similar regulatory functions. MSK1 (also designated RLPK) is a novel Rsk-related protein, which, like the p90 Rsk family members, contains two non-identical complete kinase catalytic domains.

REFERENCES

1. Alcorta, D.A., et al. 1989. Sequence and expression of chicken and mouse Rsk: homologs of *Xenopus laevis* ribosomal S6 kinase. *Mol. Cell. Biol.* 9: 3850-3859.
2. Sweet, L.J., et al. 1990. Identification of mitogen-responsive ribosomal protein S6 kinase pp90 Rsk, a homolog of *Xenopus* S6 kinase II, in chicken embryo fibroblasts. *Mol. Cell. Biol.* 10: 2413-2417.
3. Kozma, S.C., et al. 1990. Cloning of the mitogen-activated S6 kinase from rat liver reveals an enzyme of the second messenger subfamily. *Proc. Natl. Acad. Sci. USA* 87: 7365-7369.
4. Banerjee, P., et al. 1990. Molecular structure of a major Insulin/mitogen-activated 70 kDa S6 protein kinase. *Proc. Natl. Acad. Sci. USA* 87: 8550-8554.
5. Moller, D.E., Xia, C.H., Tang, W., Zhu, A.X. and Jakubowski, M. 1994. Human Rsk isoforms: cloning and characterization of tissue-specific expression. *Am. J. Physiol.* 266: C351-C359.

CHROMOSOMAL LOCATION

Genetic locus: RPS6KA1 (human) mapping to 1p36.11.

SOURCE

Rsk-1 (W-23) is an affinity purified rabbit polyclonal antibody raised against a peptide mapping within an internal region of Rsk-1 of human origin.

PRODUCT

Each vial contains 100 μ g IgG in 1.0 ml PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

Rsk-1 (W-23) is recommended for detection of Rsk-1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for Rsk-1 siRNA (h): sc-29475, Rsk-1 shRNA Plasmid (h): sc-29475-SH and Rsk-1 shRNA (h) Lentiviral Particles: sc-29475-V.


Molecular Weight of Rsk-1: 90 kDa.

Positive Controls: A-431 whole cell lysate: sc-2201, HeLa whole cell lysate: sc-2200 or MCF7 whole cell lysate: sc-2206.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2051 or ABC: sc-2018 rabbit IgG Staining Systems.

DATA



Rsk-1 (W-23): sc-130870. Western blot analysis of Rsk-1 expression in non-transfected (A) and human Rsk-1 transfected (B) 293 whole cell lysates.



Rsk-1 (W-23): sc-130870. Immunoperoxidase staining of formalin-fixed, paraffin-embedded human colon carcinoma tissue showing cytoplasmic localization.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **Rsk-1 (A-10): sc-393147** or Rsk-1 (10B1D7): **sc-81162**, our highly recommended monoclonal alternatives to Rsk-1 (W-23).