



Selenoprotein V (E-13): sc-131066

BACKGROUND

Selenium is an essential trace element that is incorporated as selenocysteine into the primary structure of selenoproteins. Nutritional deficiency of selenium decreases selenoprotein concentrations and leads to pathologic conditions. Most of the known selenoproteins are members of the glutathione peroxidase or iodothyronine deiodinase families. Selenoprotein V, also known as SELV, is a 346 amino acid testis-specific protein that contains a selenocysteine (Sec) residue at its active site and is thought to be involved in redox-related processes throughout the cell. The gene encoding Selenoprotein V maps to human chromosome 19, which is the genetic home for a number of immunoglobulin superfamily members, including the killer cell and leukocyte Ig-like receptors, a number of ICAMs, the CEACAM and PSG family and Fc receptors (Fc Rs).

REFERENCES

1. Kryukov, G.V., Kryukov, V.M. and Gladyshev, V.N. 1999. New mammalian selenocysteine-containing proteins identified with an algorithm that searches for selenocysteine insertion sequence elements. *J. Biol. Chem.* 274: 33888-33897.
2. Kryukov, G.V., Castellano, S., Novoselov, S.V., Lobanov, A.V., Zehrab, O., Guigó, R. and Gladyshev, V.N. 2003. Characterization of mammalian selenoproteomes. *Science* 300: 1439-1443.
3. Online Mendelian Inheritance in Man, OMIM™. 2003. Johns Hopkins University, Baltimore, MD. MIM Number: 607919. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Pappas, A.C., Zoidis, E., Surai, P.F. and Zervas, G. 2008. Selenoproteins and maternal nutrition. *Comp. Biochem. Physiol. B, Biochem. Mol. Biol.* 151: 361-372.
5. Zhuo, P. and Diamond, A.M. 2009. Molecular mechanisms by which selenoproteins affect cancer risk and progression. *Biochim. Biophys. Acta* E-Published.
6. Allmang, C., Wurth, L. and Krol, A. 2009. The selenium to selenoprotein pathway in eukaryotes: more molecular partners than anticipated. *Biochim. Biophys. Acta* E-Published.
7. Lescure, A., Rederstorff, M., Krol, A., Guicheney, P. and Allamand, V. 2009. Selenoprotein function and muscle disease. *Biochim. Biophys. Acta* E-Published.
8. Hatfield, D.L., Yoo, M.H., Carlson, B.A. and Gladyshev, V.N. 2009. Selenoproteins that function in cancer prevention and promotion. *Biochim. Biophys. Acta* E-Published.
9. Steinbrenner, H. and Sies, H. 2009. Protection against reactive oxygen species by selenoproteins. *Biochim. Biophys. Acta* E-Published.

CHROMOSOMAL LOCATION

Genetic locus: SELV (human) mapping to 19q13.2.

SOURCE

Selenoprotein V (E-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of Selenoprotein V of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-131066 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Selenoprotein V (E-13) is recommended for detection of Selenoprotein V of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other Selenoprotein family members .

Suitable for use as control antibody for Selenoprotein V siRNA (h): sc-97401, Selenoprotein V shRNA Plasmid (h): sc-97401-SH and Selenoprotein V shRNA (h) Lentiviral Particles: sc-97401-V.

Molecular Weight of Selenoprotein V: 37 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.