

Arp-T1 (E-12): sc-131409

BACKGROUND

Actin-related proteins are classified into Arp subclasses according to their amino acid sequence similarity to Actin. Both Arps and Actin proteins have an ATPase domain, which catalyzes the decomposition of adenosine triphosphate (ATP) into adenosine diphosphate (ADP) and a free phosphate ion to release energy. Arp-T1 (Actin-related protein T1), also known as AIP1 or ARIP1, is a 376 amino acid structural, cytoskeletal protein. Like other Arp family members, Arp-T1 contains an Actin-like ATP/ADP-binding pocket and two nuclear export signals. Although Arp-T1 shares 48% sequence identity with β -Actin, it contains several cysteine residues that are not found in other Actin family members. With significant expression in testis, Arp-T1 is implicated to play a role in spermatid formation and differentiation.

REFERENCES

1. Schafer, D.A. and Schroer, T.A. 1999. Actin-related proteins. *Annu. Rev. Cell Dev. Biol.* 15: 341-363.
2. Harata, M., Nishimori, K. and Hatta, S. 2001. Identification of two cDNAs for human Actin-related proteins (Arps) that have remarkable similarity to conventional actin. *Biochim. Biophys. Acta* 1522: 130-133.
3. Machesky, L.M. and May, R.C. 2001. Arps: Actin-related proteins. *Results Probl. Cell Differ.* 32: 213-229.
4. Heid, H., Figge, U., Winter, S., Kuhn, C., Zimbelmann, R. and Franke, W. 2002. Novel Actin-related proteins Arp-T1 and Arp-T2 as components of the cytoskeletal calyx of the mammalian sperm head. *Exp. Cell Res.* 279: 177-187.
5. Rousseaux-Prevost, R., Lecuyer, C., Drobecq, H., Sergheraert, C., Dacheux, J.L. and Rousseaux, J. 2003. Characterization of boar sperm cytoskeletal cylicin II as an Actin-binding protein. *Biochem. Biophys. Res. Commun.* 303: 182-189.
6. Online Mendelian Inheritance in Man, OMIM™. 2006. Johns Hopkins University, Baltimore, MD. MIM Number: 608535. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
7. Hara, Y., Yamagata, K., Oguchi, K. and Baba, T. 2008. Nuclear localization of profilin III-ArpM1 complex in mouse spermiogenesis. *FEBS Lett.* 582: 2998-3004.
8. LeClaire, L.L., Baumgartner, M., Iwasa, J.H., Mullins, R.D. and Barber, D.L. 2008. Phosphorylation of the Arp2/3 complex is necessary to nucleate actin filaments. *J. Cell Biol.* 182: 647-654.

CHROMOSOMAL LOCATION

Genetic locus: ACTRT1 (human) mapping to Xq25.

SOURCE

Arp-T1 (E-12) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Arp-T1 of human origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-131409 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Arp-T1 (E-12) is recommended for detection of Arp-T1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Arp-T1 (E-12) is also recommended for detection of Arp-T1 in additional species, including equine, canine and porcine.

Suitable for use as control antibody for Arp-T1 siRNA (h): sc-90996, Arp-T1 shRNA Plasmid (h): sc-90996-SH and Arp-T1 shRNA (h) Lentiviral Particles: sc-90996-V.

Molecular Weight of Arp-T1: 42 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.