

NDUFB11 (Q-13): sc-131635

BACKGROUND

Complex I (also known as NADH dehydrogenase) of the electron transport chain (respiratory chain) is an enzymatic complex that catalyzes the transfer of electrons from NADH to ubiquinone. Free energy from the reaction is conserved in the transfer of protons into the intermembrane space to create an electrochemical proton gradient, a driving force for ATP synthesis. Complex I is a complicated, multi-protein, L-shaped complex composed of at least 45 different subunits and located in the mitochondrial inner membrane. NDUFB11 (NADH dehydrogenase (ubiquinone) 1 β subcomplex subunit 11), also known as ESSS, Np15, Np17.3 (neuronal protein 17.3) or p17.3, is a hydrophobic transmembrane protein belonging to the Complex I NDUFB11 subunit family. Ubiquitously expressed, NDUFB11 localizes to the inner membrane of the mitochondrion and functions as an accessory subunit of Complex I. The cAMP-dependent phosphorylation of NDUFB11 is important for the regulation of Complex I activity.

REFERENCES

- Cui, Y., Yu, L., Gong, R., Zhang, M., Fan, Y., Yue, P. and Zhao, S. 1999. Cloning and tissue expression characterization of a full length cDNA encoding human neuronal protein p17.3. *Biochem. Genet.* 37: 175-185.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 300403. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Janssen, R.J., Nijtmans, L.G., van den Heuvel, L.P. and Smeitink, J.A. 2006. Mitochondrial complex I: structure, function and pathology. *J. Inher. Metab. Dis.* 29: 499-515.
- Petruzzella, V., Tessa, A., Torraco, A., Fattori, F., Dotti, M.T., Bruno, C., Cardaioli, E., Papa, S., Federico, A. and Santorelli, F.M. 2007. The NDUFB11 gene is not a modifier in Leber hereditary optic neuropathy. *Biochem. Biophys. Res. Commun.* 355: 181-187.
- Fernandez-Moreira, D., Ugalde, C., Smeets, R., Rodenburg, R.J., Lopez-Laso, E., Ruiz-Falco, M.L., Briones, P., Martin, M.A., Smeitink, J.A. and Arenas, J. 2007. X-linked NDUF11 gene mutations associated with mitochondrial encephalomyopathy. *Ann. Neurol.* 61: 73-83.
- Gurok, U., Bork, K., Nuber, U., Spörle, R., Nöhring, S. and Horstkorte, R. 2007. Expression of NDUFB11 encoding the neuronal protein 15.6 during neurite outgrowth and development. *Gene Expr. Patterns* 7: 370-374.
- De Rasmio, D., Panelli, D., Sardanelli, A.M. and Papa, S. 2008. cAMP-dependent protein kinase regulates the mitochondrial import of the nuclear encoded NDUF54 subunit of complex I. *Cell. Signal.* 20: 989-997.

CHROMOSOMAL LOCATION

Genetic locus: Ndufb11 (mouse) mapping to X A1.3.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

SOURCE

NDUFB11 (Q-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of NDUFB11 of mouse origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-131635 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

NDUFB11 (Q-13) is recommended for detection of NDUFB11 of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other NDUF family members.

Suitable for use as control antibody for NDUFB11 siRNA (m): sc-149878, NDUFB11 shRNA Plasmid (m): sc-149878-SH and NDUFB11 shRNA (m) Lentiviral Particles: sc-149878-V.

Molecular Weight of NDUFB11: 11 kDa.

Molecular Weight of phosphorylated NDUFB11: 18 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.