



Torsin3A (P-12): sc-131810

BACKGROUND

Torsin3A (TOR3A), also known as ATP-dependent interferon-responsive protein (ADIR), is a 397 amino acid member of the clpA/clpB family of proteins. Localized to the cytoplasm and the endoplasmic reticulum, Torsin3A is most highly expressed in stomach, lymph nodes and salivary glands. Torsin3A contains a hydrophobic N-terminal sequence, an ATP binding domain and eight putative phosphorylation sites. Highly homologous with TorsinA and TorsinB, Torsin3A is thought to be related to the AAA chaperone-like family of ATPases. Members of this family play a role in the assembly, disassembly and operation of a variety of protein complexes, confer increased tolerance to high temperature and promote specific proteolysis. Torsin3A is also referred to as an interferon responsive gene, as interaction with type 1 interferons (IFN- α and IFN- β) cause a 5- to 40-fold increase in Torsin3A mRNA levels. Three isoforms of Torsin3A exist as a result of alternative splicing events, and isoform 2 is expressed exclusively in the placenta.

REFERENCES

1. Augood, S.J., Martin, D.M., Ozelius, L.J., Breakefield, X.O., Penney, J.B. and Standaert, D.G. 1999. Distribution of the mRNAs encoding TorsinA and TorsinB in the normal adult human brain. *Ann. Neurol.* 46: 761-769.
2. Dron, M., Meritet, J.F., Dandoy-Dron, F., Meyniel, J.P., Maury, C. and Tovey, M.G. 2002. Molecular cloning of ADIR, a novel interferon responsive gene encoding a protein related to the torsins. *Genomics* 79: 315-325.
3. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 607555. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Oberlin, S.R., Konakova, M., Pulst, S. and Chesselet, M.F. 2004. Development and anatomic localization of TorsinA. *Adv. Neurol.* 94: 61-65.
5. Muraro, N.I. and Moffat, K.G. 2006. Downregulation of torp4a, encoding the *Drosophila* homologue of TorsinA, results in increased neuronal degeneration. *J. Neurobiol.* 66: 1338-1353.
6. Snider, J. and Houry, W.A. 2006. MoxR AAA+ ATPases: a novel family of molecular chaperones? *J. Struct. Biol.* 156: 200-209.
7. van Bergen, C.A., Kester, M.G., Jedema, I., Heemskerk, M.H., van Luxemburg-Heijs, S.A., Kloosterboer, F.M., Marijt, W.A., de Ru, A.H., Schaafsma, M.R., Willemze, R., van Veelen, P.A. and Falkenburg, J.H. 2007. Multiple myeloma-reactive T cells recognize an activation-induced minor histocompatibility antigen encoded by the ATP-dependent interferon-responsive (ADIR) gene. *Blood* 109: 4089-4096.

CHROMOSOMAL LOCATION

Genetic locus: TOR3A (human) mapping to 1q25.2.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

SOURCE

Torsin3A (P-12) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Torsin3A of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-131810 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

Torsin3A (P-12) is recommended for detection of Torsin3A isoforms 1, 2 and 3 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Torsin3A siRNA (h): sc-88800, Torsin3A shRNA Plasmid (h): sc-88800-SH and Torsin3A shRNA (h) Lentiviral Particles: sc-88800-V.

Molecular Weight of Torsin3A: 46 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.