

KCNH6 (S-12): sc-131963

BACKGROUND

Voltage-gated potassium channels play an essential role in controlling cellular excitability in the nervous system. They regulate a variety of properties including membrane potential as well as the frequency and structure of action potentials. KCNH6, also called potassium voltage-gated channel subfamily H member 6 or human ether-a-go-go potassium channel 2 (hEAG2), is an α subunit of a multi-pass transmembrane potassium channel family. KCNH6 functions in forming the pore of the voltage-gated channel. The channel itself is a homo- or heterotetrameric structure of pore-forming α subunits that associate with modulating β subunits. KCNH6 contains one PAS-associated C-terminal (PAC) domain, one PER-ARNT-SIM (PAS) domain and one cyclic nucleotide-binding domain. Expressed as three isoforms produced by alternative splicing, KCNH6 is present in Prolactin-secreting adenomas and throughout the brain.

REFERENCES

1. Ganetzky, B., Robertson, G.A., Wilson, G.F., Trudeau, M.C. and Titus, S.A. 1999. The EAG family of K⁺ channels in *Drosophila* and mammals. *Ann. N.Y. Acad. Sci.* 868: 356-369.
2. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 608168. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
3. Papa, M., Boscia, F., Canitano, A., Castaldo, P., Sellitti, S., Annunziato, L. and Tagliatela, M. 2003. Expression pattern of the ether-a-go-go-related (ERG) K⁺ channel-encoding genes ERG1, ERG2, and ERG3 in the adult rat central nervous system. *J. Comp. Neurol.* 466: 119-135.
4. Bauer, C.K., Wulfsen, I., Schäfer, R., Glassmeier, G., Wimmers, S., Flitsch, J., Lüdecke, D.K. and Schwarz, J.R. 2003. HERG K⁺ currents in human Prolactin-secreting adenoma cells. *Pflugers Arch.* 445: 589-600.
5. Gutman, G.A., Chandy, K.G., Grissmer, S., Lazdunski, M., McKinnon, D., Pardo, L.A., Robertson, G.A., Rudy, B., Sanguinetti, M.C., Stühmer, W. and Wang, X. 2005. International Union of Pharmacology. LIII. Nomenclature and molecular relationships of voltage-gated potassium channels. *Pharmacol. Rev.* 57: 473-508.
6. Yoshida, K. and Niwa, T. 2006. Quantitative structure-activity relationship studies on inhibition of HERG potassium channels. *J. Chem. Inf. Model.* 46: 1371-1378.
7. Ferrer, T., Rupp, J., Piper, D.R. and Tristani-Firouzi, M. 2006. The S4-S5 linker directly couples voltage sensor movement to the activation gate in the human ether-a-go-go-related gene (hERG) K⁺ channel. *J. Biol. Chem.* 281: 12858-12864.

CHROMOSOMAL LOCATION

Genetic locus: KCNH6 (human) mapping to 17q23.3; Kcnh6 (mouse) mapping to 11 E1.

SOURCE

KCNH6 (S-12) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of KCNH6 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-131963 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

KCNH6 (S-12) is recommended for detection of KCNH6 isoforms 1, 2 and 3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other KCNH family members.

KCNH6 (S-12) is also recommended for detection of KCNH6 isoforms 1, 2 and 3 in additional species, including equine and porcine.

Suitable for use as control antibody for KCNH6 siRNA (h): sc-93962, KCNH6 siRNA (m): sc-146365, KCNH6 shRNA Plasmid (h): sc-93962-SH, KCNH6 shRNA Plasmid (m): sc-146365-SH, KCNH6 shRNA (h) Lentiviral Particles: sc-93962-V and KCNH6 shRNA (m) Lentiviral Particles: sc-146365-V.

Molecular Weight of KCNH6: 110 kDa.

Positive Controls: Hep G2 cell lysate: sc-2227 or PC-12 cell lysate: sc-2250.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.



Try **KCNH6 (46): sc-135959**, our highly recommended monoclonal alternative to KCNH6 (S-12).