

# RNF40 (E-13): sc-132077

## BACKGROUND

Ubiquitination is an important mechanism through which three classes of enzymes act in concert to target short-lived or abnormal proteins for destruction. The three classes of enzymes involved in ubiquitination are the ubiquitin-activating enzymes (E1s), the ubiquitin-conjugating enzymes (E2s) and the ubiquitin-protein ligases (E3s). RNF40 (RING finger protein 40), also known as BRE1B, Staring or RBP95, is a 1,001 amino acid nuclear protein that contains one RING-type zinc finger. Expressed ubiquitously with highest expression in heart, testis and pancreas, RNF40 functions as an E3 ubiquitin-protein ligase that regulates the monoubiquitination and subsequent degradation of select residues on target proteins, such as Histone H2B and Syntaxin 1. In addition, RNF40 forms a ubiquitin ligase complex with UBCH6 (an E2 enzyme) and together, these proteins play a crucial role in regulation of the histone code. Four isoforms of RNF40 exist due to alternative splicing events.

## REFERENCES

1. Wen, H., et al. 2000. RBP95, a novel leucine zipper protein, binds to the retinoblastoma protein. *Biochem. Biophys. Res. Commun.* 275: 141-148.
2. Li, D., et al. 2002. The role of adenovirus-mediated retinoblastoma 94 in the treatment of head and neck cancer. *Cancer Res.* 62: 4637-4644.
3. Chin, L.S., et al. 2002. Staring, a novel E3 ubiquitin-protein ligase that targets Syntaxin 1 for degradation. *J. Biol. Chem.* 277: 35071-35079.
4. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 607700. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Hwang, W.W., et al. 2003. A conserved RING finger protein required for Histone H2B monoubiquitination and cell size control. *Mol. Cell* 11: 261-266.
6. Wood, A., et al. 2003. BRE1, an E3 ubiquitin ligase required for recruitment and substrate selection of Rad6 at a promoter. *Mol. Cell* 11: 267-274.
7. Zhu, B., et al. 2005. Monoubiquitination of human Histone H2B: the factors involved and their roles in HOX gene regulation. *Mol. Cell* 20: 601-611.

## CHROMOSOMAL LOCATION

Genetic locus: RNF40 (human) mapping to 16p11.2; Rnf40 (mouse) mapping to 7 F3.

## SOURCE

RNF40 (E-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of RNF40 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-132077 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-132077 X, 200 µg/0.1 ml.

## APPLICATIONS

RNF40 (E-13) is recommended for detection of RNF40 isoforms 1, 2, 3 and 4 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with other RNF family members.

Suitable for use as control antibody for RNF40 siRNA (h): sc-93054, RNF40 siRNA (m): sc-153050, RNF40 shRNA Plasmid (h): sc-93054-SH, RNF40 shRNA Plasmid (m): sc-153050-SH, RNF40 shRNA (h) Lentiviral Particles: sc-93054-V and RNF40 shRNA (m) Lentiviral Particles: sc-153050-V.

RNF40 (E-13) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of RNF40: 95 kDa.

Positive Controls: A-673 cell lysate: sc-2414 or HeLa whole cell lysate: sc-2200.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



RNF40 (E-13): sc-132077. Western blot analysis of RNF40 expression in HCT 116 whole cell lysate.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.